

A
COMPENDIOUS SYSTEM
OF
Natural Philosophy.
With NOTES

Containing the
MATHEMATICAL DEMONSTRATIONS,
AND
Some Occasional REMARKS.

PART II.
HYDROSTATICS and PNEUMATICS.
To which are annexed some DISSERTATIONS
relating to these Subjects.

By J. ROWNING, M. A.
Rector of ANDERBY in LINCOLNSHIRE, and late
Fellow of MAGDALEN COLLEGE in CAMBRIDGE.

The THIRD EDITION.

L O N D O N:
Printed for SAM. HARDING Bookseller, on the
Pavement in St. Martin's Lane.
M.DCC.XXXVII.

N

Of

I.

a ft

*
Bodi
all su
Fluid
the M

A
COMPENDIOUS SYSTEM
OF
Natural Philosophy.

PART II.
HYDROSTATICS.

CHAP. I.

Of the Phænomena which arise from the mutual Action of the Particles of Fluids upon one another.

I. **I**N the former Part of this Essay, I have laid down, and explained the general *Laws of Nature*, and from thence deduced those Phænomena, which are in a strict and proper Sense * denominated *Me-*

* In a larger Sense all the Effects and Operations of natural Bodies upon one another may be called Mechanical ; as being all subject to the general Laws of Motion. In Hydrostatics Fluids are governed by the Laws of Mechanism, as much as the Mechanical Powers themselves are ; the same holds of the

chanical; I proceed now to an Explanation of such, as Philosophers have comprehended under the Name of *Hydrostatics*; the Intention of which is to explain the Nature of *Fluids*, and the Manner, wherein they act upon one another, and upon Solids.

II. The Nature of a *Fluid**, as distinguished from that of a *Solid*, or hard Body, consists in this, *viz.* that its Particles are so loosely connected together, that they readily move out of their Places, when pressed with the least Force one Way more than another †.

Rays of Light, as will be seen when we come to Optics; and in the larger Bodies of the Planetary System, Mechanism equally prevails, as has been demonstrated by Sir *Isaac Newton*; which we shall endeavour to make out (when we treat of Astronomy) so far as the Nature of our Design will permit.

* Some Philosophers make the following Distinction in Fluids; those which flow or spread themselves till their Surfaces become level or horizontal, they call *Liquid*; in Contradistinction to Flame, Smoak, Vapour, &c. which are also Fluids, but do not acquire such a Surface. Those, which are capable of exciting in us the Idea of Moistness, as Water, &c. they call *Humid*, distinguishing them thereby from Air, Quicksilver, and melted Metals. But those Distinctions are quite unnecessary in a Philosophical Sense: the Surfaces of all Fluids being level, when not prevented by the Bodies about them; and Humidity is only a *relative* Quality; for tho' Quicksilver will not moisten, or stick to a Man's Fingers, it will to Silver or Gold.

† The common Definition, *Fluidum est cujus partes impressioni cuicunque cedunt, & cedendo facillime moventur inter se*, though it expresses very well the Nature of a *compressible* Fluid, as Air, yet does not correspond to that of Water, whose Parts have been found to yield to no Force, with which they have been compressed, unless it was greater on one Side than on the other. The Definition therefore seems imperfect, as not expressing that

Inequa-

From whence Philosophers conclude, that they are exceedingly minute, smooth, and round *; it being otherwise impossible they

Inequality of Pressure, which is requisite to move the Parts of some Fluids one among another.

* It is commonly observed that the Roundness of the Particles conduces very much to Fluidity, not only as it disposes them to move one among another with greater Facility, but because round Bodies touching one another in few Points only, the Force with which they mutually attract each other, is the weaker. But, upon this Supposition, the Particles of a Fluid ought to move with less Freedom one among another, by how much the greater the Weight is, with which they are compressed, (for it is the same Thing in this respect, whether they press against each other by Virtue of their own Attraction, or by some external Force) but of this we have no Experience. A Diver, upon plunging out of his Bell at the Bottom of the Sea, never finds the Water less fluid, notwithstanding the great Pressure from above. Mr. Boyle having caused a Tadpole to be put into a Vessel of Water, and to be pressed with a very great Force, tells us, that in Appearance it found no Inconvenience from thence, but swam about with the same Freedom and Briskness as ever.

Quære, Whether the Particles of which Fluids consist, are in Contact with each other, or not? Perhaps, they are prevented from approaching, nearer than to a certain Distance, by a repelling Power, diffused around each single Particle. The Observation, that Water is not render'd less fluid by Pressure, seems to favour this Opinion; and the Property, which the Air has of expanding or contracting itself, according to the Weight which it sustains (as shall be shewn Chap. 3.) proves beyond Contradiction, that its Particles are endued with such a Power. But then if the Particles of all Fluids have this Power, it will follow that they ought to be in some Measure capable of being reduced into less Space by Pressure, as Air is; which they have not as yet appeared to be. Further, since it has been proved (Part I. Chap. 3.) that if the Parts of Fluids are placed just beyond their natural Distances from each other, they will approach and run together; and if placed farther asunder still, will repel each other; it follows, upon the foregoing Supposition,

should move with such Freedom one among another, upon the least Inequality of Pressure.

tion that each Particle of a Fluid must be surrounded with three Spheres of Attraction and Repulsion one within another : the innermost of which is a Sphere of Repulsion, which keeps them from approaching into Contact ; the next, a Sphere of Attraction diffused around this of Repulsion, and beginning where this ends, by which the Particles are disposed to run together into Drops ; the outermost of all, a Sphere of Repulsion, where by they repel each other, when removed out of that Attraction.

Now, if this *Hypothesis* should be found to be true ; and we might agreeably to the above-mentioned Consequence of it, suppose, that the Particles of all Bodies attract and repel each other *alternately* at different Distances, perhaps we might be able to solve a great many Phenomena relating to small Bodies, which now lie beyond the reach of our Philosophy. However, upon the Supposition of the three Spheres of Attraction and Repulsion just mentioned, nothing is more easy, than to see how Solids may be converted into Fluids, and Fluids into Solids (as is done in Liquefaction and Freezing) ; for allowing, that the first or innermost Sphere of Repulsion is capable, like that of the Particles of Air, of being augmented by Heat, and diminished or totally suspended by Cold ; it follows, that Bodies must be more or less fluid, in Proportion to the Degree in which they are affected by Heat or Cold : for when the Action of the first Sphere of Repulsion is diminished or destroyed by Cold, the Particles of the Fluid must necessarily be brought into closer Contact with each other by the Force of the circumambient Attraction, and by that means constitute an harder Body than before. But, we must not dwell too much upon an Hypothesis which wants Proof ; I shall only add, that altho' some Fluids, as Water, have not been *as yet* contracted in their Dimensions, or made to take up less Space than they naturally do, by any Force with which they have been compressed by *Art* ; yet there are none but are *naturally* contracted by Cold, from whence it seems reasonable to infer, that their Particles are at least *capable* of being brought into closer Contact, which is some Confirmation of this Doctrine.

It is an obvious Objection to this, that Water by freezing is augmented in its Bulk ; but this may be owing to those Bubbles or Vacuities, observable in the Water after it is frozen, which

III. Those Particles, considered separately, are endued with all the common Properties of Matter, and subject to the same Laws of Motion and Gravitation with larger Bodies. To enquire therefore into the Nature of Fluids, is to consider, what Appearances a Collection of very small round Bodies, subject to those Laws, will exhibit under different Circumstances. In order to which, it is usual with Hydrostatical Writers to consider a Fluid, as divided into several perpendicular Columns contiguous to each other. Sometimes it is convenient to conceive it divided into thin Plates or *Strata* lying upon one another. In some Cases, the same Fluid is considered, as distinguished both these Ways, *viz.* into perpendicular Columns, and also into thin Strata or Plates. Figure 1. represents a Vessel filled with a Fluid to the Height EF, and divided into the Columns GH, IK, LM, &c. and also into the Strata RS, TV, XY, &c.

IV. From this Observation, concerning the Properties of the Particles considered separately, immediately results the following Proposition, *viz.* that in a Vessel, whose Form is such as is represented by ABCD, (*Fig 1.*) the Quantity of Pressure, which each Stratum su-

which were not in it before ; and not to any general and uniform Removal of the Particles of the Fluid from each other, which the Objection, if it is of any Force against what has been advanced, must suppose.

stains,

stains from the Weight of the incumbent Fluid, is in Proportion to the Number of those Strata, which rest upon it, that is, as the Height of the Surface of the Fluid above it; for if we suppose the Strata of equal Thickness, the Quantity sustained is proportionable to the Number of Strata of which it consists.

V. When the Surface of a Fluid is horizontal or level, each Particle thereof is disposed to continue in its Place, being sustained therein by the contiguous ones.

Let the Fluid be supposed to be divided into Strata, each of the Thickness of a Particle of the Fluid; and if the Truth of this Proposition be denied, let the Particle *mm* be one of those which is not sustained in its Place by the contiguous ones, but is moving from thence towards some other Part of the Vessel, *e.g.* towards D. Now, since all the other Particles of that Stratum are at an equal Depth below the Surface of the Fluid with this, they also sustain an equal Degree of Pressure (by the last Proposition,) consequently for the same Reason that one of them is moving towards D, the rest may all be said to be moving in the same Direction: But this cannot be true of the whole Stratum, while the Vessel is entire, and therefore of none of its Parts. Now, the like Reasoning will hold against the Motion of the Particle *mm* towards any other Part of the Vessel; from whence it follows,

follows, that each Particle of the Fluid, is sustained in its Place by the contiguous ones, and therefore disposed to continue at Rest *.

VI From hence is derived a fundamental Proposition in Hydrostatics, *viz.* That when the Surface of a Fluid is level, whatever Pressure any single Particle or small Portion of it sustains from the contiguous ones on one Part, it sustains the same on all the rest; that is, it is pressed by them with an equal Degree of Force on all Sides †.

For, by the Definition of the Fluid (§. 2.) each Particle is disposed to give Way, and move out of its Place, when the Pressure is not equal on all Sides; and (§. 5.) each Particle is pressed by the contiguous ones in such a Manner, that it is sustained in its Place thereby; it is therefore pressed with an equal Degree of Force on all Sides.

* This shews us the Absurdity of some Philosophers, who make the Nature of a Fluid, as contradistinguished from that of a Solid, to consist solely in the intestine Motion of its Particles.

† This Proposition with its Corollary is not strictly speaking true, unless the Particle or Portion of Fluid we speak of is supposed void of Gravity, for it presses downwards with a Force equal to the Weight of those Particles which rest upon it, added to its own; whereas the Force with which it presses upwards is only equal to the Weight it sustains, *viz.* that of the incumbent Fluid. But the Particles of Fluids are so exceedingly minute, and the Gravity of each so very small, that the Error arising from hence can never be sensible.

Corol.

Corol. From hence it follows, that each Particle or small Portion of a Fluid presses with the same Degree of Force in all Directions on those which are contiguous to it. For, by the third Law of Nature, every Particle presses upon the contiguous ones with the same Degree of Force, with which it is pressed upon by them.

VII. The Surface of a Fluid becomes level by its own Gravity, when no external Force prevents it from being so.

For, the Particles of Fluids press in all Directions with Forces proportionable to the Height of their Surfaces (Cor. §. 6. and §. 4.) if then the Surface be not level, the different Parts of the same inferior Stratum will be pressed not only downwards, but sideways against each other with unequal Forces; the greater Pressure therefore overcoming the weaker, the Particles which sustain the least Pressure, will be driven out of their Places and raised up, till the Surface becomes level; the Surface being level, each Particle will be equally pressed in every Direction, (§. 8.) all therefore will remain at Rest, and the Surface continue in that State *

* This Demonstration, as also the foregoing, is founded upon a Supposition, that Bodies tend downwards by their Gravity in Lines parallel to each other, which though *physically* true, is not strictly so, their Tendency being towards the Center of the Earth, and consequently in Lines which meet in a Point: and therefore, if we would be accurate, the Fluid contained

ta
an
th
fel
the
and
of t
sho
libr
equa
the
posi
Spac
is div
be of
EF is
tion o
is the
Consec
convex
Convex
This
Earth's
Portions
be negle
Mariner
Building
Manner,
Mast and
from their

VIII. Fluids gravitate in Fluids of the same Kind.

This Proposition is a necessary Consequence of what has been observed about the Nature of the Particles of which Fluids consist, *viz.* that they are solid, and endued with the same Properties with other Bodies. The Reason why their Gravity is not sensible in the Fluid, is because the lower Parts sustain

tained in a Vessel should be considered, as divided into Columns and Strata, as represented Figure the second, where ABD is the Earth, C its Center, EFGH a Fluid contained in a Vessel, and divided into Columns, which if continued down to the Center of the Earth, would there terminate in a Point C; and into the concentric Strata, *ab, cd, &c.* having the Center of the Earth for the Center of their Convexity. And then we should find that the Particles of the Fluid will not be in *Æquilibrium* with each other, till all the Parts of its Surface are at equal Distances from the Center of the Earth, forming thereby the Surface EF, concentric to that of the Earth. For, supposing the Fluid EFGH continued down to C, so as to fill the Space ECF; it is evident the Columns, into which the Fluid is divided, cannot be of equal Lengths, and therefore cannot be of an exact Counterpoise to each other, unless the Surface EF is a Portion of a Sphere, whose Center is C: but the Action of the Parts of the Fluid upon each other at the Surface, is the same, whether the lower Part GCH be a Fluid, or not. Consequently the Surfaces of Fluids are not level or plain, but convex, having the Center of the Earth for the Center of their Convexity.

This Convexity, by Reason of the great Distance of the Earth's Center, approaches so near to a Plane, that in small Portions of it, the Difference is not sensible, and therefore may be neglected: but at Sea 'tis evident to Sense; for when the Mariners put to Sea, the Shore first disappears, then the lower Buildings, afterwards the Towers, Mountains, &c.; in like Manner, when they approach a distant Ship, the Top of its Mast and Sails appear first, while the Ship itself is intercepted from their View, by the Convexity of the Water between them.

the upper, and hinder them from descending. But, it does not follow from thence, that the Gravity of the uppermost is entirely taken away, as some * Philosophers have imagined; for, as much as the lower Parts press against the upper in sustaining them; just so much do the upper by their Reaction press upon the lower. Thus, the Weight of the whole Fluid is communicated to the Vessel, which therefore weighs according to the Quantity it contains, notwithstanding the Pressure of each Portion of the Fluid, taken separately, seems not to effect it.

IX. The Pressure of a Fluid is in Proportion to its perpendicular Height, and the Quantity of Surface against which it presses,

This Proposition admits of four Cases.

1. When the Fluid is contained in a Vessel of the same Dimension from Top to Bottom, and held in an erect Position, as that represented Fig. 1. it is evident, the Pressure of the Fluid upon the Bottom will be in Proportion to its Magnitude, and the perpendicular Height of the Surface of the Fluid above it. For, conceiving it divided into Columns, the Pressure upon the Bottom, by the fourth Proposition, will be as the Length or Height of the Column: and it will also be as the

* This was the Notion of the *Cartesians*, who held, that when a Fluid is mixed with another of the same Kind, it loses its own Weight thereby.

Number of them, because the Quantity of Fluid, which presses upon the Bottom, is in that Proportion, that is, as the Magnitude of the Bottom pressed upon. But when the Vessel is inclined or irregular, the Truth of this Proposition is so far from being evident, that it has been commonly looked upon as a Paradox.

2. Let the Vessel ABCD (*Fig. 3.*) be filled with a Fluid to the Height EF, and held in an inclined Position, as there represented; I say, the Pressure of the Fluid is proportionable to the Magnitude of the Base CD, and FG or HD the perpendicular Height of the Surface of the Fluid above it.

For supposing the Fluid divided into the Strata EI, KM, LO, &c. so far, as the first Stratum EI is prevented from pressing upon KI the Surface of the next inferior Stratum, by being in some Measure supported by the Side of the Vessel FI, so far is its Pressure augmented by the Reaction of the opposite Side EK upon it, which is exactly equal to the Action of the former, because the Fluid pressing every Way alike, at the same Depths below the Surface, exerts an equal Force against both these Sides. The Surface therefore of the second Stratum is pressed with the same Degree of Force with which it would be, if the Quantity of Fluid contained in the former Stratum was included within the Space

HKQI, which is exactly equal to it, as having the same Base KI and the same perpendicular Height QI *. Now, this being true of each Stratum, their Pressure upon CD the Base of the Vessel is the same, as if they were all placed perpendicularly over it, and filled the Space RHCD; which they would do, since the Sum of their perpendicular Heights QI, KS, LT and NV are equal to HD the perpendicular Height of this Space, and each of their Bases KI, LM, &c. is equal to CD its Base †. But, by the foregoing Case, if the Space RHCD was filled with a Fluid, the Pressure of it would be proportionable to the Dimensions of the Base CD, and the perpendicular Height DH, therefore it is the same in the inclined Tube ABCD ||.

3. Let the Vessel ABC be irregular, as represented *Figure* the fourth, and filled with

* 31. *El.* II:

† 31. *El.* II.

|| Perhaps it may be thought more Geometrical, to demonstrate this Proposition with the Generality of Authors from the Property of the inclined Plane. They consider AD the lower Side of the Tube, as an inclined Plane, on which the Fluid contained within it rests, and argue that it loses thereby a Part of its Weight in Proportion to the Length of the Plane, and therefore occasions no greater Pressure upon the Base, than if the Vessel was held erect, and filled only to the same perpendicular Height, as when inclined. But this Demonstration proves too much, for by this Way of Reasoning, one might shew, that the Pressure of the Fluid EFCD upon the Base CD is less than the Pressure of RHCD a Column of the same Fluid having the same Base and equal perpendicular Height with it. For both the inclined, and the perpendicular Column contain the

a Fluid to the Height D, I say, the Pressure of the Fluid upon the Base C, is proportionable to the Magnitude of the Base, and CD the perpendicular Height of the Surface of the Fluid above it.

In order that the Proof of this Proposition may be the better understood, we must premise the following *Lemma*. viz,

That when a Fluid passes through a Pipe, as AB, (*Fig. 5.*) which in some Parts is larger than in others, the Moment, or Force with which it moves, is every where the same. For while the Fluid is passing through the Tube, its Velocity in every Part will be reciprocally as the Quantity of Matter; for Instance, it will be as much greater at C, than it is at D, as the Quantity passing through C at any Instant of Time is less than at D, and so of the rest; because a less Quantity would be conveyed through the smaller Parts of the Tube in the same Time, unless it should move faster there in Proportion to the smallness of them. Now the Momentum of Bodies is partly owing to the Quantity of Matter, and partly to the Velocity; (as explained Part I. Chap. 9. §. 1.) consequently what the Fluid, which is actual-

the same Quantity of Fluid, upon Account of the Equality of their Bases, and perpendicular Heights; but that rests upon an inclined Plane which this does not, and therefore presses less upon the Base. But this is contrary both to Demonstration and Experience, the Argument therefore proves too much.

ly

ly passing through the narrower Parts of the Tube, wants in Quantity, is compensated by its Velocity in those Parts, and what it wants in Point of Velocity in the other Parts is made up by the Quantity passing through them; so that the Moment is the same in every Part of the Tube, whether larger or narrower *. The same is true, whatever be the Position the Tube is held in.

Let us now conceive the Fluid in the Vessel ABC (*Fig. 4.*) to be distinguished into the Strata EF, GH, IK, &c. Let us also imagine the Bottom of the Vessel C to be moveable, that is, capable of sliding up and down the narrow Part of the Vessel, *e. g.* from C to GH, (without letting any of the Fluid run out). Let it further be supposed that this moveable Bottom is drawn up or let down with a given Velocity, while the Vessel itself is fixed and immoveable; it is evident the lowermost Stratum, which is contiguous to the Bottom, will be raised or let down with the same Velocity, and will thereby have a Moment proportionable to that Velocity and the Quantity of Matter it contains: But by the Lemma, all the rest of the

* Thus, we may observe in a River or Canal, that by how much the Breadth or Depth is less in any Part, so much the more rapid is the Stream in that Part; and on the contrary, where it is wider and deeper, the Motion of the Water is more gentle and languid. So that the Moment, with which it flows, is the same in every Part.

t II.

the
d by
ants
s is
ough
n e-
r or
r be

the
l in-
also
be
and
v. g.
the
posed
r let
Vef-
evi-
nti-
let
ere-
hat
con-
the

how
n the
rary,
ter is
hich
rata

Fig. 1. P. 7.




Fig. 2. P. 11.




Fig. 3. P. 13.




Fig. 4. P. 14.




Fig. 5. P. 15.





Cha

Stra
quer
(tha
if th
Part
Mon
been
Pres
with
tom
Stra
Flu
Stra

A

6.)
sure
to t
the

I
as b
give
tum
mob
of a
Ves
it is
is p
and
is th

*

throu

Strata will have the same Moment, consequently the Moment of all taken together, (that is, of the whole Fluid,) is the same, as if the Vessel had been no larger in any one Part, than it is at the Bottom, (for then the Moment of each Stratum would also have been as great as that of the lowermost); the Pressure therefore, or Action of the Fluid, with which it endeavours to force the Bottom out of its Place, is as the Number of Strata, that is, the perpendicular Height of the Fluid, and the Magnitude of the lowermost Stratum, that is, the Bottom.

Again, suppose the Vessel ABC (*Fig. 6.*) fill'd with a Fluid to D; I say the Pressure upon the Bottom BC, is proportionable to the Dimensions of the Bottom, and to DE the perpendicular Height of the Fluid.

For, if we suppose the Bottom moveable, as before, and raised up or let down with a given Velocity, the Moment of every Stratum will be the same with that of the lowermost, by the Lemma; therefore the Moment of all taken together, is the same, as if the Vessel had been no less in any one Part, than it is at the Bottom; consequently the Pressure is proportionable to the perpendicular Height, and the Magnitude of the Bottom. This Case is the Converse of the former *.

* Upon this is founded the Practice of conveying Water through Pipes from Place to Place, &c. For from hence it follows,

From hence it follows, that, if a Vessel is made of such a Form, as is represented (*Fig. 7.*) by ABCDEFG, and filled with a Fluid to the Height C, the Weight which the Bottom sustains, is as great as it would be, had the Vessel been IKFG (which is every where of the same Dimensions, that the other is of at the Bottom) and filled to the Top IK. Because the Pressure, by the Proposition, is proportionable to the Bottom and perpendi-

follows, that if one End of a Pipe is laid in a Reservoir of Water, the Fluid will run into the Pipe, till it rises to a Level at the other End with its Surface in the Reservoir. Thus let ABC (*Fig. 8.*) represent a Reservoir or Basin of Water, DGE a Pipe laid from thence to E. If E the End of the Pipe is placed above the Line ABF, the Level of the Water in the Reservoir, the Water will run into the Pipe, till it rises in the other End to F the Level with AB; at which Time the Water in the Pipe will be in *Æquilibrium* with that in the Reservoir, and remain at Rest. But if the End of the Pipe is below the Surface of the Water in the Reservoir, it will continue to run out, till they are reduced to a Level. For, let GH be the lowest Part of the Pipe, then since F the perpendicular Height of the Fluid on one Side, is equal to B the perpendicular Height of the Fluid on the other, and GH, which (being the Place where the Fluids press one against another,) may be consider'd as a Base to both, is common; it follows from this Proposition, that the Pressures on each Side are exactly equal; and therefore being in contrary Directions will necessarily destroy each other, and the Fluid will remain in *Æquilibrium*. But while the End E is below the Level, this *Æquilibrium* cannot be obtained; and therefore the Fluid will continue to run out.

For the same Reason, when two or more Tubes communicate with each other, the Surface of the Fluid they contain will stand at the same Level in all.

cular

cular Height, which in both Cases are the same *.

X. The Velocity, with which a Fluid spouts out at an Hole in the Bottom of a Vessel, is equal to that, which a Body would acquire by falling freely from the Level of the Surface of the Fluid to the Hole.

Let there be a large cylindrical Tube ABCD (*Fig. 9.*) in the upper Part of which let us imagine a Cylinder of Ice FGHI ex-

* The Reason why the Vessel ABCDEFG, with the Fluid contained in it, does not weigh so much, as the Vessel IKFG, when full to the same Height, notwithstanding the Pressure upon the Bottom is the same in both, is, because ABDE the upper Part, or Cover of the former Vessel, is pressed upwards by the Fluid below it, with a Force equal to the Endeavour the Fluid in the small Tube BCD has to descend. Which Endeavour is the same that it would be, if the Tube BCD comprehended also the two Spaces ICBA and CKED, its Moment being the same in both Cases by the Demonstration; the Cover therefore is pressed upwards with a Force equal to the Weight of as much Fluid as would fill the two Spaces ICBA and CKED; consequently the Vessel, whose Form is ABCDEFG, is so much lighter than the other, that is, as much as the Fluid it contains is less.

From hence arises this Paradox, that the least Quantity of Fluid may be made to raise any Weight how great soever it be.

For since the Cover ABDE is pressed upwards with a Force equal to the Weight of as much Fluid, as would fill the two Spaces ICBA and CKED, and those Spaces may be enlarged at Pleasure in Height, by lengthening the Tube BCD (which at the same Time must be made proportionably smaller, otherwise the same Quantity of Fluid will not fill it;) it follows, that the same Quantity of Fluid may be made to press the Cover upwards with a given Force; if that Cover then is made moveable, any Weight that is laid upon it may be supported thereby.

C

actly

actly fitting it; let it further be supposed, that HI, the lower Surface of the Ice, is continually melting, so as to afford a Stream of Water running down the Middle of the Tube. Now the Form of this Stream of Water will necessarily be such, as is represented in the Figure by HLI; for the Water falling freely will descend faster and faster like other Bodies, causing thereby the Stream to become narrower and narrower. Now, let it be supposed, that the Tube has a Bottom as CD, with an Hole in it at K, just sufficient to let the Stream pass freely; it is evident, there will be no Obstruction on this Account, but that the Fluid will pass through the Hole with such Velocity, as it naturally acquires by falling from HI, the lower Surface of the Ice. And if we suppose M and N, the empty Parts of the Tube, to be filled with Water, the Water will press equally upon the Sides of the Stream in every Direction (§. 6.) and therefore will be no Impediment to its Motion on that Account. Lastly, let us suppose the Ice taken away, and the Stream supplied from the Water at the Sides, as is the Case when a Fluid runs out through the Bottom of a Vessel; then will the Velocity, with which the Water flows through the Hole, continue the same; for so far as the Water coming from the Sides, endeavours to descend itself, so far it obstructs the Descent of the Stream, and no farther;
and

and consequently causes no Alteration in the Velocity or Quantity of Fluid running out. The Velocity therefore, with which the Fluid passes through the Hole, is equal to that, which a Body would acquire by falling freely from the Level of the Surface of the Fluid to that Place.

If the Hole is made in the Side of the Vessel at the same Distance below the Surface, the Velocity will be the same, on Account of that equal Tendency Fluids have to move every Way alike*.

XI. The Velocity, with which a Fluid spouts out from the Bottom or Side of a Vef-

* Upon this Principle is founded the Practice of making artificial Fountains. For if to a Vessel or Reservoir ABCD, (Fig. 10.) filled with a Fluid to the Height EF, be fixed the Pipe CH, with a small Aperture at K, the Fluid will spout up from thence to FL, the Level of the Surface of the Fluid in the Vessel. For, by this Proposition, it will spout from K with such a Velocity, as a Body would acquire by falling from FL the Level of the Surface to the Aperture at K, that is, such as will carry it from the Aperture to the Level; because that Velocity which a Body acquires by falling from a certain Height, is sufficient to carry it back to the same Height from whence it fell.

But in Practice the Height the Fluid rises to, is less than that of the Level of its Surface in the Reservoir: this is owing to the Resistance it meets with from the Air, its Friction against the Sides of the Pipe, &c. It is found impossible to make it much exceed the Height of an hundred Feet: For, when it spouts out of the Aperture with a Velocity necessary to carry it higher, the Stream is immediately dashed to Pieces by the Resistance of the Air; whereby it loses its Force, and is prevented from rising to any considerable Height.

fel, is as the square Root of the Height of its Surface above the Hole *.

The Cause, why a Fluid spouts out through an Hole made in the Bottom or Side of a Vessel, is the Pressure or Weight of the Fluid incumbent upon the Hole; from whence it should seem, that the Velocity ought to be as the Pressure; but if so, then the Quantity run out would also be as the Pressure (for the faster the Liquor flows, the greater is the Quantity thrown out in a given Time, and *vice versâ*) consequently, upon this Supposition we should have two Effects, each depending on the same Cause, and equal to it, which is absurd. 'Tis not then the Quantity of Fluid run out, nor the Velocity with which it flows, but its Moment, or both these multiplied together, (Part I. Ch. 9. §. 1.) that is the true and adequate Effect of the Pressure. Now these being ever in the same Ratio, with each other, 'tis necessary, in Order that

* This Proposition may be otherwise demonstrated from the last, in the following Manner. For, since the Velocity with which a Fluid spouts out through an Hole in the Bottom or Side of a Vessel, is equal to that which a Body would acquire by falling from the Level of the Surface of the Fluid to the Hole, and the Velocities Bodies acquire by falling are as the square Roots of the Heights they fall from (Part I. Chap 5. §. 5.) it follows, that the Velocity, with which a Fluid spouts out from an Hole in the Bottom or Side of a Vessel, is as the square Root of the Height of the Level of the Surface of the Fluid above the Hole.

the

the Effect may be proportionable to its Cause, that each of them be only as the square Root of the Pressure: For then, being multiplied together, their Product, or the Moment of the spouting Fluid, is adequately as the Pressure which occasions it; but the Pressure is as the perpendicular Height (§. 4.) therefore the Velocity, and also the Quantity of Fluid spouting out, is only as the square Root of the Height of its Surface above the Hole.

To give an Instance or two; suppose two Holes made in the Side of a Vessel, the one an Inch below the Surface of the Fluid it contains, the other four Inches; the Velocity with which the Liquor flows out of the lower Hole, will not be four Times as great, as that with which it flows through the upper, notwithstanding the Pressure is four Times greater: For if it should, the Quantity run out in a given Time would also be four Times greater, consequently the Effect produced would be sixteen Times greater than it is at the upper Hole, that is, four Times greater than the Cause; which is absurd. Whereas the Velocity and Quantity of Matter will each be only twice as great as they are above, producing thereby a Force or Moment only four Times as great, which is proportionable to the Cause. So, if an Hole were made sixteen Times lower than the first, the Velocity and Quantity of Matter will not be each sixteen Times greater than at the other,

other, but only four Times greater apiece, and so the Moment sixteen Times greater, as the Pressure is *.

XII. When a Current of Water, or other Fluid, falls perpendicularly upon the Surface

* From hence, we may see the Error some of the foreign Mathematicians have fallen into with Regard to the Forces of moving Bodies, who contend that they are as the Squares of the Velocities multiplied by the Quantities of Matter. For, from this Proposition it is, that one of the principal Arguments brought in Favour of this Opinion is derived. They argue thus, *Effects* are ever proportionable to their *Causes*, the *Pressure* of the incumbent Fluid is the *Cause* of its spouting out, the *Force* with which it spouts out, is the *Effect*; but by this Proposition the *Pressure* is as the *Square* of the *Velocity* it flows with, therefore the *Force* is likewise as the *Square* of the *Velocity*. True, it is so; but let us see the Consequence. The *Force*, with which the Fluid spouts out is not only owing to the *Velocity*, but the *Quantity* run out in a given *Time*. They have each their Share in producing the *Force*, consequently the *Force* is in a *Ratio* compounded of both, or as the *Product* of one multiplied by the other, or, which comes to the same Thing, (since as was observed before, they are in the same *Ratio* with each other) as the *Square* of either of them. From hence it is that the *Forces* of *Fluids* in *Motion* are said to be as the *Squares* of their *Velocities*; not that they are so in *Virtue* of those *Velocities*, as such, but in *Virtue* of them, and the *Quantities* of *Matter* taken together, or because the *Squares* of the *Velocities* is the same Thing with the simple *Velocities* multiplied by the *Quantities* of *Matter*. Therefore when it is said, the *Forces* of *Fluids* are as the *Squares* of the *Velocities*, that Part of the *Force* which arises from the *Quantity* of *Matter* is really taken into Consideration. How ridiculous then must it be in those Gentlemen to fetch an Argument from hence to prove, that the *Forces* of *Bodies* in *Motion* are as the *Squares* of the *Velocities* and *Quantities* of *Matter* too, when they are as the *Squares* of the *Velocities*, only because the *Quantities* of *Matter* are implied in them?

II.

and
the

her
ace

eign
rces
ares
tter.
Ar-
hey
uses,
out-
et ;
the
the
the
out
run
pro-
om-
by
was
her)
the
ares
nose
nti-
the
ities
n it
Ve-
man-
ri-
an
dies
nti-
Ve-
in

of

Fig. 6. P. 17.




Fig. 7. P. 18.




Fig. 8. P. 17.





Fig. 9. P. 20.



Fig. 10. P. 21.





CH

of
aga
Fo
to
wh
He
it
ty

let
fill
EF
the
in
of
ing
the
In

in
fusp
to
the
port
whi
tom
itfel
ed

*
the P
Water
moirs

of a Plane, or flows against it, (as the Wind against the Sail of a Ship, or the like) the Force, which the Fluid exerts upon it, is equal to the Weight of a Column of the same Fluid, whose Base is equal to the Plane, and its Height such, that a Body falling freely through it would acquire the same Degree of Velocity with which the Fluid moves*.

In Order to demonstrate this Proposition, let us suppose the Vessel ABCD (*Fig. 11.*) filled with a Fluid, and having a large Hole EF in the Bottom, then will the Pressure of the Fluid cause a Stream to flow out, which in the Hole itself will have such a Degree of Velocity, as a Body would acquire by falling freely from the Surface of the Fluid in the Vessel to the Hole (as demonstrated §. 10.) In the midst of this Hole, and consequently in the Stream, let us suppose a Plane as PQ suspended, but somewhat less than is sufficient to fill the Hole, lest it stop the Current of the Water. Now, 'tis certain, this Plane supports a Column of the Fluid, equal to that which presses upon any other Part of the Bottom of the Vessel of equal Dimensions with itself (for being thus placed, it may be looked upon as a Part of the Bottom) but every

* From this Proposition is deduced the Method of computing the Power of a Machine, which is to be moved by Wind or Water, &c. See an Instance of such a Calculation in the Memoirs of the Royal Academy of Sciences for the Year 1725.

Part bears a Column, whose Base is equal to its own Dimensions, and its Height the same with that of the Surface of the Fluid in the Vessel: Consequently this Plane supports such a Column, that is, it is resisted by the Stream with a Force equal to the Weight of a Column, whose Base has the same Dimensions with itself, and whose Height is equal to that of the Surface of the Fluid in the Vessel, that is, such an Height as a Body by falling freely from, would acquire a Velocity equal to that with which the Fluid moves.

XIII. The Pressure of a Fluid against a perpendicular Bank or Sluice, &c. is equal to the Weight of a Column of the same Fluid, whose Base is equal to so much of the Bank as is below its Surface, and which has half the Depth of the Fluid for its Height *.

If the Pressure upon every Part of the Bank from the Surface to the Bottom, was as great as it is at the Bottom, the Pressure against it would be equal to the Weight of a Column, whose Base is equal to so much of the Bank as is under the Surface of the Fluid, and which has the *whole* Depth of the Fluid for

* From hence we see the Reason, why the Water of the Sea or great Lakes is as easily kept within their Banks (setting aside the Force which arises from the Motion of the Waves, &c.) as that of the narrowest Canal, viz. because the Pressure of Fluids is not in Proportion to their Surfaces, but their Depths.

its Height ; for the Pressure upon every Part of the Bank at the Bottom is equal to the Weight of a Column, whose Base corresponds to the Part pressed upon, and whose Height is that of the Depth of the Fluid ; consequently if the Pressure was the same every where from Top to Bottom, it would be equal to the Weight of as many such Columns as would answer to all the Parts of the Bank : But the Pressure every where diminishes in Proportion as we approach the Top, where it is Nothing ; it is therefore but *half* * what it would be in the other Case ; from whence the Proposition is clear.

C H A P. II

Of the Effects Fluids have on Solids immersed therein.

I. **T**HE *Specific Gravity* of a Body is that, by which it is said to be heavier, or lighter than another of a different Kind : Thus Lead is said to be specifically heavier than Cork ; because supposing an equal Bulk of each, the one would be heavier than the other.

* Because the Sum of a Number of Terms in Arithmetical Progression beginning from Nothing, is half the Sum of an equal Number of Terms, each of which is equal to the last in the Progression.

D

From

From hence it follows, that a Body, specifically heavier than another, is also more dense, that is, contains a greater Quantity of Matter under the same Bulk, because Bodies weigh in Proportion to the Quantities of Matter they contain (Part I. Chap. 3. §. 7.)

II. If a Solid be immersed in a Fluid of the same specific Gravity with itself, it will remain suspended therein, in whatever Part of the Fluid it is put.

Let the Body FGHI (*Fig. 12.*) be immersed in the Fluid ABCD to the Depth MN, or any other whatever; I say, it will continue in the same Part of the Fluid, when left to itself, without either rising towards the Surface, or sinking towards the Bottom.

For, the Body being (by the Supposition) of equal Gravity with the Fluid, the Weight of the Column KLHI, which consists partly of Fluid, and partly of the Body, is the same as if it had been all Fluid; consequently HI, that Part of the Surface of the Stratum MN, which lies immediately under the Body, is pressed with the same Degree of Force, that any other Part of the same Dimensions is, and therefore the whole Column KLHI will be supported in its Place. Now, the same being true of the Column KLHI, whatever be its Length, 'tis evident the Body will be suspended in its Place at any Depth.

III. But, if the Body is specifically heavier than the Fluid, in which it is immersed, it will subside to the Bottom: For then, in whatever Part of the Fluid it is put, the Column KLHI will always be heavier than an equal Column, that consists all of Fluid; consequently HI, that Part of the Stratum MN which lies immediately under the Body, will suffer a greater Pressure, than any other Part of the same Dimensions; and therefore will give way, and permit the Body to subside continually, till it reaches the Bottom.

IV. On the contrary, if the Body is specifically lighter than the Fluid, it will rise to the Top, in what Part of the Fluid soever it is put. For then, the Column KLHI will always be lighter than an equal Column which is all Fluid; consequently HI will be less pressed downwards, than any other Part of the same Stratum of equal Dimensions, and will therefore continually rise up, carrying the Body with it, till it arrives at the Top.

V. A Body, being laid on the Surface of a Fluid specifically heavier than itself, sinks into it, till the immersed Part takes up the Place of a Quantity of Fluid, whose Weight is equal to that of the whole Body.

Let EFGH (*Fig. 13.*) be a Body, floating on a Liquor specifically heavier than itself, it will sink into it, till the immersed Part IKGH takes up the Place of so much Fluid

as is equal to it in Weight. For, in that Case, GH, that Part of the Surface of the Stratum upon which the Body rests, is pressed with the same Degree of Force, as it would be, was the Space IKGH full of the Fluid; that is, all the Parts of that Stratum are pressed alike, and therefore the Body, after having sunk so far into the Fluid, is in *Æquilibrio* with it, and will remain at Rest.

From hence it follows, that a Body is as much specifically lighter, than the Fluid on which it floats, as the immersed Part is less than the Whole. For, by how much the less the immersed Part is, so much the less Fluid is equal in Weight to the whole Body; that is, the Body is so much the lighter in Respect of the Fluid. And, if the same Body be made to float successively in Fluids, whose specific Gravities differ among themselves, (but all exceed that of the Body), the lighter the Fluids are, so much greater will be the Part immersed*.

VI. A Body, suspended in a Fluid specifically lighter than itself, loses a Part of its Weight (or rather communicates it to the

* This Phenomenon is what gave rise to the Hydrometer, an Instrument of great Use in ascertaining the Genuineness of Liquors; for it rarely happens, that the adulterated and the genuine Liquor (however they may agree in Appearance) are of the same specific Gravity.

Fluid) equal to that of a Quantity of Fluid of the same Bulk.

Let us, instead of supposing the Body suspended in the Fluid, imagine it to be away, and its Place filled with the Fluid; now 'tis evident, this being of the same specific Gravity with the circumjacent Fluid, will be entirely supported by it, or, if we suppose the Body to be of the same specific Gravity with the Fluid, it will be wholly suspended by it; we see therefore the Pressure of the circumambient Fluid, whereby it endeavours to buoy up the Body, is equivalent to the Weight of so much Fluid, as would fill the Place the Body takes up. But, since the Fluid presses only on the Surface of the Body, that Pressure is the same, whatever be the specific Gravity of the Body; the Body therefore loses so much of its Weight, as the Fluid would naturally buoy up; that is, so much, as is the Weight of a Quantity of Fluid of the same Bulk.

This Proposition affords us a Method of determining the Relation, which the specific Gravities of Bodies, whether Fluid or Solid, bear to each other. For, whereas by weighing a Solid in a Fluid specifically lighter than itself, we find the absolute Weight of a Quantity of the Fluid equal to it in Bulk (*viz.* the Weight the Solid loses) the Relation, that Weight bears to the Weight of the Solid, is the Relation of their specific Gravities; because

cause the Weights of Bodies, whose Bulks are equal, are as their specific Gravities: consequently, if the same Solid is weighed successively in different Fluids (all lighter than itself) we gain the Relation, which the specific Gravity of each bears to that of the Solid, and therefore to one another. Again, if different Solids are weighed in the same Fluid, the Relation, which the specific Gravity of that Fluid bears to each Solid is had, and therefore also the specific Gravities of the Solids among themselves*.

* Upon this is founded the Use of the Hydrostatical Balance for determining the specific Gravities both of Solids and Fluids. The Practice is thus. First, let the Solid be weighed in Air, that is, out of the Fluid; afterwards in it (this ought to be done by suspending it at one End of the Balance by a String, that is as nearly of the same specific Gravity with the Fluid made Use of as possible, and letting it sink into the Fluid till it is wholly immersed below the Surface; if the Fluid is Water, an Horse-Hair is most convenient to hang the Body at the End of the Balance by) then subtract its Weight in the Fluid from its Weight in Air, the Difference is what it loses in the Fluid. This done, say, by the Rule of Proportion, as the Weight lost in the Fluid is to its Weight in Air, so is Unity, or any Number taken at Pleasure, to a Fourth, which by its Relation to the former, will express the Relation of specific Gravity of the Solid to that of the Fluid. Thus, the Relation, which the specific Gravity of the same Fluid bears to that of various Solids, or of the same Solid to that of various Fluids, and consequently the Relation of the specific Gravities of all among themselves may be obtained.

C H A P. III.

Of the Air.

I. **T**HAT Part of Natural Philosophy, which treats of the Properties of the Air, and the Effects of its *Pressure* and *Elasticity*, is called *Pneumatics*.

II. Air is a thin transparent elastic Fluid surrounding the Earth to a certain Height, and taken all together, is called the *Atmosphere*.

III. That Air is a Fluid, is evident from the easy Passage it affords to Bodies moving in it: For this shews it to be a Body, whose Parts easily yield to a Pressure, that is greater on one Side than on the other, which is the Definition of a Fluid.

IV. Air gravitates towards the Earth, or is heavy like other Bodies.

To prove this, we have Abundance of Arguments both from Sense and Experiment. Thus, when the Hand is applied to the Orifice of a Vessel, it readily perceives the Weight of the incumbent Atmosphere, as soon as the Air included in the Vessel begins to be drawn out. Thus, Glass-Vessels exhausted of their Air (if not strong enough to sustain the Pressure of the incumbent Atmosphere) are crushed to Pieces by the Weight of the Air without

out. When the Air is exhausted out of a Vessel, the Vessel weighs less than before. With a great many more Experiments generally mentioned by Authors on this Subject *.

V. The exact Weight of the incumbent Air is determined by filling a Tube with Mercury, and immersing the open End in a Vessel of the same Fluid: For then the Mercury will run down the Tube, till its Surface is fallen only to the perpendicular Height of about twenty nine or thirty Inches above the Surface of the Mercury in the Vessel; if the same Experiment is made with Water, the Surface of it will stand at about the Height of thirty two Feet above the Surface of that in the Vessel; the Column of Mercury in one Case, and the Column of Water in the other, exactly balancing the Weight of a Column of Air, which reaches to the Top of the Atmosphere, and presses upon the Surface of the Fluid in the Vessels. This is what is called the Torricellian Experiment, from TORRICELLI the Inventor, and is the same with the common Barometer.

From hence it follows (Chap. I. §. 9.) that all Bodies, at the Surface of the Earth, sustain as great a Weight from the Pressure of the Air, as is that of a Column of Water,

* See Boyle's Tracts, or *Gravesande Lib. II. P. III.*

whose

whose Height is thirty two Feet, and its Base equal to the Surface of the Body pressed upon *.

VI. That the Suspension of the Mercury in the Barometer depends on the Pressure of the external Air, is beyond all doubt; for if the Barometer is included in the Air-Pump, the Mercury falls in the Tube, in Proportion as the Air is exhausted out of the Receiver; and if the Air is let in again gradually, the Mercury reascends proportionably, till it reaches its former Height.

VII. That the Atmosphere is extended to a determinate Height, appears from hence; *viz.* that when the Torricellian Tube is removed to a more elevated Place, the suspended Column of Mercury becomes shorter; which is, because a shorter Column of Air presses upon it; or, that the Tube, in this Situation, is nearer the Top of the Atmosphere.

* The Pressure of the Atmosphere upon every square Inch, near the Surface of the Earth, is about fifteen Pounds, being equal to the Weight of a Column of Mercury, whose Height is thirty Inches and its Base one square Inch. For, such a Column of Mercury would weigh about fifteen Pounds. The Weight of the Atmosphere therefore, which presses upon a Man's Body, is equal to so many Times fifteen Pound, as the Surface of his Body contains square Inches.

The Reason, why a Person suffers no Inconvenience from so great a Pressure, is owing to the Air included within the Pores and Fluids of the Body, which by its Reaction is a Counterpoise to the Pressure of the external Air.

VIII. The Elasticity of the Air is that Property by which it contracts itself into less Space, when an additional Pressure is laid upon its; and recovers its former Dimensions, when the Pressure is taken off. This is accounted its distinguishing Property, all the rest being common to it with other Fluids.

Of this we have numerous Proofs. Thus, a Bladder full of Air being compressed by the Hand, the included Air gives way; but when the Pressure is taken off, the Air expands itself, and readily fills up the Cavity, or Impression made in the Surface of the Bladder. And, if a larger Quantity of Air, than is naturally pressed into a Vessel by the Weight of the incumbent Atmosphere, is forced into it by the *Condenser* (an Engine for that purpose) and if that Air is afterwards let out by opening the Vessel, the Remainder is found to be of the same Weight as at first; from whence it follows, that the Air, by means of its Elasticity or Spring, drives out all that which was forced in by the Condenser, recovers its former Dimensions, and fills the Vessel as before.

IX. From hence, together with what has been observed about the Pressure of the Atmosphere, it follows, that the Air near the Surface of the Earth, is compressed into a much narrower Space by the Weight of the Air above, than that which it would naturally take up, was it free from that Pressure;

sure; accordingly it is found by Experiment, that, when the Pressure of the Atmosphere is taken off from any Portion of the Air, it immediately expands itself into a vast Extent. Hence it is, that thin Glass Bubbles or Bladders filled with Air, being included in the Receiver of the Air-Pump, are broke in Pieces by the Spring of the Air, which they contain within them, when the Pressure of the external Air is taken off. Thus a Bladder quite flaccid, containing only a small Quantity of Air in it, swells upon the Removal of the external Air, and appears distended, as if it contained as great a Quantity as possible. The same Effect is found in carrying a Bladder somewhat flaccid to a more elevated Place, for there the external Pressure being less, the Air included in the Bladder is in some Measure free from the Pressure of the Atmosphere; it therefore dilates itself, and distends the Bladder as in the former Case.

X. It is found by Experiment, that the Quantity of Space into which Air may be contracted by Pressure, is reciprocally proportionable to the compressing Force. From whence it follows, that the Density of the Air is proportionable to the Pressure which it sustains; because the less the Space is, into which a given Quantity of it is contracted, the denser it is. As to the utmost Degrees of Expansion and Contraction, which the Air is capable

ble of, they are as yet unknown. In several Experiments made by Mr. BOYLE, Air in its natural State, that is, pressed only with the Weight of the incumbent Atmosphere, dilated itself, when that Pressure was taken off, into more than thirteen thousand Times the Space it took up before. And, he was able so far to compress it, that it should take up more than five hundred and twenty thousand Times less Space than that, into which it would dilate itself, when free from its Pressure *.

XI. From this Property it follows, that the Air in the inferior Parts of the Atmosphere is more dense, than that, which is at great Heights in the same; or, that the Density of the Air decreases continually, as we approach the Top of the Atmosphere. For the Density of the Air is proportionable to the Force

* See Boyle's Tracts and Experiments on the Spring and Pressure of the Air.

Various have been the Opinions of Philosophers concerning the Cause of this prodigious Spring in the Particles of Air; some holding it to depend on their Figure, which they suppose to resemble in some Manner little Bundles of Twigs or the Branches of Trees; some think them like Fleeces of Wool, others conceive them as rolled up like Hoops, or the Springs of Watches, and endeavouring to expand themselves by Virtue of their Texture. But Sir Isaac Newton is of Opinion, that such a Texture is by no Means sufficient to account for that vast Power of Expansion observed above; but that each Particle is endued with a repelling Force which encreases as they approach one another, and accordingly keeps them asunder at Distances reciprocally proportionable to the Pressure they sustain.

See Hales's Statical Essays. Vol. I. Chap. 6.

with

with which it is compressed, and that Force continually decreases, as we approach the Top.

Was the Density of the Atmosphere every where the same, that is near the Surface of the Earth, its Height (as is computed from the Quantity of Pressure it exerts in raising the Mercury in the Barometer) would be about five Miles. But whereas its Density continually decreases, as we approach the Top, and it is uncertain how far the Particles may expand themselves, where there is little or no Pressure, the true Height cannot be obtained. It is computed to continue of a sensible Density to the Height of about forty five, or fifty Miles.

XII. The Elasticity of the Air produces the same Effects with its Pressure.

For, Action being equal to Reaction, the Force, which the Spring of the Air exerts in endeavouring to expand itself, is equal to the Force with which it is compressed; just as it is in the Spring of a Watch, which exerts no Force, but in Proportion as it is wound up; consequently a Quantity of Air in such a State of Contraction, as it would be compressed into by the Weight of the incumbent Atmosphere, exerts a Force equal to that Weight. If a Quantity of Air therefore is included in a Vessel, and is of the same Density with the circumambient Air, its Pressure against the Sides of the Vessel is equal to the Weight of the Atmosphere. Thus, Mercury is sustained
to

to the same Height by the elastic Force of Air included in a Glass Vessel, no Way communicating with the external Air, as by the Weight of the Atmosphere itself.

XIII. The Elasticity of the Air is augmented by Heat and diminished by Cold *. For if a Bladder, which is about half filled with Air, is laid before the Fire, it will, when it is sufficiently heated, be distended and burst. Thus, Glass Bubbles being laid upon the Fire immediately burst with great Violence by the augmented Spring of the included Air.

XIV. The Density of the Air thus continually varying, according to the different Degrees of Heat and Cold, to which it is exposed, makes it difficult to ascertain its true specific Gravity. RICCIOLUS estimates it to be to that

* This Property is found in all Bodies both Solid and Fluid. but in a much less Degree, than it is in Air, Thus, if a Flask be filled with Water, only to the lower Part of the Neck, and is then set upon the Fire, the Water, when it begins to grow warm, will rise into the Neck, and continue to ascend, as the Heat is increased. And when a Wire or Bar of Iron is heated, it is augmented both in Length and Diameter.

Upon this Property depends the Phænomena of the Thermometer, which is a Glass Bubble with a small hollow Stem arising from it. This Bubble and Part of the Stem is usually filled with Mercury, or Spirit of Wine, which will rise or fall in the Stem, as they are affected by the Heat or Cold of the external Air. If a sufficient Degree of Heat is suddenly applied to this Instrument, the Liquor is observed to descend a little before it rises, because the Glass distending itself, the Capacity of the Bubble is augmented, before the included Liquor is affected by the Heat.

Chap. 4. *The Resistance of Fluids* 41

of Water, as one to a Thousand: MERSENNUS, as one to one Thousand three Hundred: Mr. BOYLE, by more accurate Experiments, found it to be, as one to nine Hundred and Thirty eight; and thinks, that, all Things consider'd, the Proportion of one to a Thousand may be taken as a Medium; for there is no fixing any precise Proportion, because not only the specific Gravity of Air, but that of Water also, is continually varying. However, by some Experiments made since with more Accuracy before the *Royal Society*, the Proportion has been fixed at about one to eight Hundred and Eighty.

XV. Air is necessary for the Preservation of Animal and Vegetable Life; neither will Fire subsist without it. The Reason of this is as yet unknown to Philosophers. Mr. HALES by several curious Experiments in his Statical Essays makes it probable, that 'tis owing to its Elasticity*.

C H A P. IV.

Of the Resistance of Fluids.

THE Resistance a Body meets with in moving through a Fluid, is of three Kinds. The first arises from the Friction of the Body against the Particles of the Fluid;

* See his *Analysis* of it. Statical Essays, Vol. I. Chap. 6.

42 *The Resistance of Fluids.* Part II.

the second from their Cohesion or Tenacity among themselves; the third, from their Inactivity, or the Tendency they have in common with other Bodies, to keep the Places they possess.

The first, *viz.* that which arises from the Friction of the Body against the Particles of the Fluid, is very inconsiderable; for whatever the Weight is, which presses the Particles of a Fluid together, the Freedom, with which a Body moves through it, is not sensibly diminished thereby. As was observed Chap. I. §. 2. in the Notes.

The second, or that which arises from the Tenacity of the Particles of the Fluid, is as the Time the Body takes up in passing thro' it^{*}; for the shorter the Time is in which the

* We have a very curious Argument in Confirmation of this, and which at the same Time illustrates the Manner in which a Body makes its Way through a tenacious Fluid, by Sir *Isaac Newton* himself, in a Postscript to a Letter in the Philosophical Transactions No 371. It is as follows. ' Suppose Pieces of
' fine Silk, or the like thin Substance, extended in parallel
' Planes, and fixed at small Distances from each other, Suppose
' then a Globe to strike perpendicularly against the Middle of
' the outermost of the Silks, and by breaking through them to
' lose Part of its Motion. If the Pieces of Silk be of equal
' Strength, the same Degree of Force will be required to break
' each of them; but the Time, in which each Piece of Silk re-
' sists, will be so much shorter as the Globe is swifter; and the
' Loss of Motion in the Globe consequent upon its breaking
' through each Silk, and surmounting the Resistance thereof,
' will be proportional to the Time in which the Silk opposes it-
' self to the Globe's Motion; insomuch that the Globe by the
' Resistance of any one Piece of Silk, will lose so much less of
' its Motion as it is swifter. But on the other Hand, by how
' much

Chap. 4. *The Resistance of Fluids.* 43

Force of Cohesion is broke through, the less Effect it has in resisting the Motion of the Body. This Species of Resistance is also very small, except in glutinous and viscid Fluids, whose Parts are not easily separated.

The third Species is the principal Resistance, that Fluids give to Bodies, and arises from their Inactivity or the Tendency the Particles, of which they consist, have to continue at Rest. The Quantity of this Resistance depends on the Velocity the Body moves with on a double Account: For in the first Place, the Number of Particles put into Motion by the moving Body in any determinate Space of Time, is proportionable to the Velocity wherewith the Body moves; and in the next Place, the Velocity with which each of them is moved, is also pro-

' much swifter the Globe moves, so many more of the Silks it
' will break through in a given Space of Time; whence the
' Number of the Silks, which oppose themselves to the Mo-
' tion of the Globe in a given Time, being reciprocally pro-
' portional to the Effect of each Silk upon the Globe, the Re-
' sistance made to the Globe by these Silks, or the Loss of
' Motion the Globe undergoes by them in a given Time, will
' be always the same.

' Now if the Tenacity of the Parts of Fluids observes the
' same Rule, as the Cohesion of the Parts of these Silks; name-
' ly, that a certain Degree of Force is required to separate and
' disunite the adhering Particles, the Resistance arising from the
' Tenacity of Fluids must observe the same Rule, as the Re-
' sistance of the Silks; and therefore in a given Time the Loss
' of Motion, a Body undergoes in a Fluid by the Tenacity of
' its Parts will in all Degrees of Velocity be the same; or in
' fewer Words, that Part of the Resistance of Fluids, which
' arises from the Cohesion of their Parts, will be uniform.

portionable to the Velocity of the Body ; this Species therefore of Resistance is in a duplicate Proportion, or as the Square of the Velocity, with which the Body moves through the Fluid *.

II. Farther the Resistance a Body moving in a Fluid meets with from thence, may be considered with Regard to the Fluid ; and then it will be found to be more or less, according to the Density of the Fluid. For by how much denser the Fluid is, so much the greater Number of Particles are to be put into Motion by the Body, in Order to make its Way through it.

III. The next Thing to be considered is the Effects of the Resistance of Fluids upon Bodies moving in them ; that is, the Retardation, which they cause in the Motion of a Body by their Resistance, or the Quantity of Motion they destroy.

* This may be otherwise demonstrated from the twelfth Section of the first Chapter ; for from thence it follows, that the Resistance a Fluid gives to a Solid against which it moves, is proportionable to the Height a Body must fall from, to acquire such a Degree of Velocity as the Fluid moves with : But the Heights Bodies fall from are as the Squares of the Velocities they acquire by falling ; consequently the Resistance a Fluid gives to a Solid, against which it moves, is also as the Square of its Velocity. Now it matters not, as to the Resistance, whether the Fluid moves against the Solid, or whether it be at Rest, and the Solid moves in it ; the Resistance therefore which a Fluid gives to a Solid moving in it, is as the Square of the Velocity, with which it moves.

And

Chap. 4. *The Resistance of Fluids.* 45

And this, in similar Bodies of equal Magnitudes, is inversely as their Densities, or the Quantity of Matter they contain; for by how much the greater the Quantity of Matter in any Body is, so much the more easily does it overcome the Resistance it meets with from the Fluid. Thus, we see the Resistance of the Air has a much less Effect in destroying the Motion of an heavy Body, than of a light one which has the same Dimensions.

IV. In similar Bodies of equal Densities, but different Magnitudes, the Retardation is inversely as their homologous Sides. For the Resistance Bodies meet with in a Fluid, is inversely as the Quantities of Matter they contain (by the last), that is inversely as the Cubes of their homologous Sides; and it is also directly as their Surfaces, because 'tis by them that they move the Fluid out of its Place; that is, directly as the Squares of their homologous Sides; consequently the Retardation is inversely as their homologous Sides *.

Having given the fundamental Principles of *Hydrostatics*, and shewn how Fluids, both compressible and incompressible, are disposed to act upon each other, and upon Solids by their Pressure, Motion, Elasticity, and Resist-

* Because the inverse Ratio of the Cubes of any Numbers being compounded with the direct Ratio of the Squares of the same, gives the inverse Ratio of the Numbers themselves.

46 *The Resistance of Fluids.* Part II.

ance ; I proceed now to account for some of the more remarkable Phænomena of Nature, in which they are in Part or altogether concerned : And this I design for the Subject of the following Dissertations.



D I S-

DISSERTATION I.

Of Sound.

WHEN the Parts of an elastic Body are put into a tremulous Motion by Percussion or the like ; so long as the Tremors continue, so long is the Air, included in the Pores of that Body, and likewise that which presses upon its Surface, affected with the like Tremors and Agitations : Now, the Particles of Air being so far compressed together by the Weight of the incumbent Atmosphere, as their repulsive Forces permit, (as has been explained Chap. 3.) it follows, that those, which are immediately agitated by the reciprocal Motions of the Particles of the elastic Body, will, in their Approach towards those which lie next them, impel them also towards each other ; and thereby cause them to be more condensed, than they were by the Weight of the incumbent Atmosphere, and in their Return will suffer them to expand themselves again ; whereby the like Tremors and Agitations will be propagated to the next ; and so on, till having arrived at a certain Distance from the Body, the Vibrations cease, being gradually destroyed by a continual successive Propagation of Motion to fresh Particles of Air throughout their Progress.

Thus

Thus it is that Sound is communicated from a tremulous Body to the Organ of Hearing. Each Vibration of the Particles of the sounding Body is successively propagated to the Particles of the Air, till it reaches those which are contiguous to the *Tympanum* of the Ear, (a fine Membrane distended across it,) and these Particles in performing their Vibrations impinge upon the Tympanum, which agitates the Air included within it; and that being put into a like tremulous Motion, affects the auditory Nerve, and thus excites in the Mind the Sensation or Idea of what we call *Sound*.

Now since the repulsive Force of each Particle of Air is equally diffused around it every Way; it follows, that when any one approaches a Number of others, it not only repels those which lie before it, in a right Line; but all the rest, laterally according to their respective Situations: that is, it makes them recede every Way from itself, as from a Center: and, this being true of every Particle, it follows, that the aforesaid Tremors will be propagated from the sounding Body in all Directions, as from a Center: And farther, if they are confined for some Time from spreading themselves by passing through a Tube or the like, will, when they have passed through it, spread themselves from the End in every Direction. In like Manner, those which pass through an Hole in an Obstacle they meet with

with in their Way, will afterwards spread themselves from thence, as if that was the Place where they began; so that the Sound, which passes through an Hole in a Wall or the like, is heard in any Situation whatever, that is not at too great a Distance from it. Something analogous to this we may observe in the Motion of Waves upon the Surface of a Fluid, which are propagated equally through all Parts of the Surface in a Circle, though occasioned not by a circular, but reciprocal Motion and Agitation of the Finger or any other Body in a straight Line.

Since the repulsive Force, with which the Particles of Air act upon each other, is reciprocally as their Distances; (Chap. 3. §. 10.) it follows, that when any Particle is removed out of its Place by the Tremors of a sounding Body, or the Vibrations of those which are contiguous to it, it will be driven back again by the repulsive Force of those towards which it is impelled, with a Velocity proportionable to the Distance from its proper Place; because the Velocity will be as the repelling Force. The Consequence of this is, that, let the Distance be great or small, it will return to its Place in the same Time; (for the Time a Body takes up in moving from Place to Place will always be the same, so long as the Velocity it moves with is proportionable to the Distance between the Places.) The Time therefore

fore in which each Vibration of the Air is performed, depends on the Degree of Repulsion in its Particles : And so long as that is not altered, will be the same at all Distances from the tremulous Body ; consequently, as the Motion of Sound is owing to the successive Propagation of the Tremors of a sounding Body through the Air, and as that Propagation depends on the Time each Tremor is performed in, it follows, that the Velocity of Sound varies as the Elasticity of the Air, but continues the same at all Distances from the sounding Body.

And as the Velocity, with which Sound is propagated, depends on the Elasticity of the Air, so it does also on its Density ; for when the Density of the Air is augmented, while its Elasticity remains the same *, a great Num.

* Perhaps it will not appear to every one, how the Density of the Air can be augmented without a proportional Increase of its Elasticity ; because *ceteris paribus*, the nearer the Particles approach each other, the stronger is the Action of their repulsive Force.

But it is to be considered, that when the Air becomes colder, its Elasticity is diminished, and then the Particles are brought closer together by the Pressure of the Atmosphere, till they acquire an Elasticity equal to what they had before ; viz. such as answers to the Pressure they sustain (Chap. 3 §. 2.) From hence we may infer, that the Propagation of Sound is slower in Winter than in Summer, when the Mercury in the Barometer is at the same Height ; for the Pressure of the Air being the same, its Elasticity which depends upon it, is so too ; but the Air is denser by Reason of the Cold, and therefore its Vibrations slower.

ber of Particles will move forwards and backwards in each Vibration; now, since we suppose the Cause by which they put each other into Motion (*viz.* their Elasticity,) the same, they will each receive a less Degree of Velocity; and so the Vibrations will be performed in a longer Time, whence the Succession of them will be slower, and the Progress of the Sound proportionably retarded*.

Whereas the undulatory Motion of the Air, which constitutes Sound, is propagated in all Directions from the sounding Body; it will frequently happen, that the Air, in performing its Vibrations, will impinge against various Objects, which will reflect it back, and

* The Method of determining the Velocity with which Sound is propagated, is (by the Help of a short Pendulum) to estimate the Time, which passes between seeing the Fire of a Gun at a Distance, and hearing the Report. Its great Velocity makes it difficult to be determined exactly; accordingly Authors differ much in their Accounts. The most accurate Observers Dr. *Halley* and Dr. *Durham* have found it to be about one Thousand one Hundred and Forty two Feet, which is almost a Quarter of a Mile, in a Second.

The usual Experiments to prove that the Air is necessary for the Propagation of Sound, are such as these. A small Bell, being put into the Receiver of the Air-Pump, may be heard at a considerable Distance, before the Air is exhausted out of it; but when the Air is much rarified by exhausting, can scarcely be heard at all. When the Air is condensed, the Sound is augmented in Proportion to the Condensation. These Experiments do not only succeed in forced Rarefactions and Condensations, but in such also as are Natural; Sound being observed to be much weaker on the Tops of high Mountains, where the Air is less condensed by the Weight of the incumbent Atmosphere, than in the Valleys below.

so cause new Vibrations the contrary Way ; now, if the Objects are so situated, as to reflect a sufficient Number of Vibrations back (*viz.* such as proceed different Ways) to the same Place, the Sound will be there repeated, and is called an *Echo* *. And, the greater the Distance of the Objects is, the longer will be the Time, before the Repetition is heard. And when the Sound in its Progress meets with Objects, at different Distances, sufficient to produce an Echo, the same Sound will be repeated several Times successively, according to the different Distances of those Objects from the sounding Body ; which makes what is called a *repeated Echo*.

If the Vibrations of the tremulous Body are propagated through a long Tube, they will be continually reverberated from the Sides of the Tube into its Axis, and by that means prevented from spreading, till they get out of it ; whereby they will be exceedingly encreased and the Sound rendered much louder than it would otherwise be †.

* In *Woodstock Park* near *Oxford*, there is an Echo, which repeats distinctly seventeen Syllables by Day, and twenty by Night. See *Plot's Natural History of Oxfordshire*.

The Reason, why it repeats more Syllables by Night, than by Day, is because the Air, being colder at that Time, is more dense ; and therefore the Return of the first Vibrations being slower, gives Time for the Repetition of more Syllables.

† This is the Case in the *Stentorophonic Tube* or *Speaking-Trumpet*.

The Difference of *Musical* Tones depends on the different Number of Vibrations communicated to the Air, in a given Time, by the Tremors of the sounding Body; and the quicker the Succession of the Vibrations is, the acuter is the Tone, and *è contra*.

A musical Chord performs all its Vibrations, whether great or small in the same Time. For, if a String is stretched between two Pins, and a Force is applied to the middle Point, to draw it out of its rectilinear Situation; it is found by Experiment, that the Distance (if it be small) to which it is drawn, is as the Force applied; consequently the Velocity, with which it returns, when left to itself, will be as the Space it has to move over; it will therefore perform all its Vibrations in the same Time: This is the Reason why the same Chord,

Trumpet. See *Kircher de Re Musica. Lib. 9. Par. 4. Ode Philos. Natur. Princip. p. 293.*

Upon this Principle it is, that Sound is conveyed from one Side of a *Whispering-Gallery* to the opposite one, without being perceived by those who stand in the Middle. The Form of a *Whispering-Gallery* is that of a Segment of a Sphere, or the like arched Figure; and the Progress of the Sound through it may be illustrated in the following Manner

Let ABC (*Fig. 14.*) represent the Segment of a Sphere; and suppose a low Voice uttered at D, the Vibrations expanding themselves every Way, some will impinge upon the Points E, E, &c. from thence be reflected to the Points F, from thence to G, and so on, till they all meet in C; and by their Union there cause a much stronger Sound, than in any other Part of the Segment whatever, even at D the Point from whence they came.

however struck, produces the same Note. It is also found by Experiment, that when Strings of equal Diameters, but different Lengths, are equally stretched, the longer they are, so much the less Weights will draw them from their rectilineal Situation to the same Distance; the Forces therefore by which they return are less, and the Times of their Vibrations longer.

When two Chords preform their Vibrations in equal Times, the Tone produced is called an *Unison*. If one performs two, while the other one; 'tis an *Octave*. If one three, while the other two; 'tis a *Fifth*. If one three, while the other four; 'tis called a *Fourth*, &c.

To make an Unison Sound, it is not necessary, that the Vibrations of the two Strings should actually concur, but only that they should be performed in equal Times; so that they would always concur, if they began at the same Instant. For the Ear perceives not the single Vibrations distinctly, but only finds that Difference, which proceeds from the Intervals of Time, that pass between them *.

* Upon these Principles we may account for that remarkable Phænomenon in *Musick*, that an intense Sound being raised, either with the Voice or a sonorous Body, another sonorous Body near it and in Unison with it, will thereby be made to sound. For the Vibrations of the Air, which correspond to the Tremors of the first sounding Body, agreeing exactly, in Point of Time, with those, which are capable of being given to the
other

other Body at Unison with it; when they have by their first Impulse communicated a small Degree of Motion to it, will, by conspiring with it as it moves forwards and backwards, continually increase its Motion, till it becomes sensible. The contrary happens in Strings which are in Dissonance with each other; for in this Case, though the first Vibration of one may give Motion to the other, yet their Vibrations not being performed in equal Times, the second will come *unseasonably*, i. e. when the other is moving the contrary Way, and obstruct its Motion. It is farther observable, that in two Strings, one of which vibrates twice, while the other once; if the first be sounded, the two Extrems of the other will each sound an Unison with it, while the middle Point remains at Rest. So if one vibrates thrice, while the other once, the last will be divided into three Parts each of which will sound an Unison with it, and the two Points between those Parts will remain at Rest. For otherwise that which vibrates twice, while the other once, must necessarily interfere with it at every second Vibration; and that which vibrates thrice, while the other once, would interfere with it at every third; so that it would not be put into a sufficient Motion to produce a Sound. But, when it is divided by the quiescent Points, it becomes so many Strings at Unison with the former, each of which easily receives its Vibrations from thence.

From hence likewise it is, that, if we take two or three drinking Glasses, and put some Water or other Fluid into each of them, and place them near to each other, taking Care to fill them, to such Heights, that (when struck) their Tones shall be in Unison; and then, if we slide the Finger along the Brim of one of the Glasses, pressing pretty strongly upon it, (which will cause it to sound) we shall see the Surface of the Fluids in the other Glasses begin to tremble; which shews that the Vibrations of the first Glass cause the like in the other at Unison with it; though not perhaps in a Degree sufficient to produce a Sound, strong enough to be heard distinctly from the former.

Thus it is that some Persons are able to break a drinking Glass by a Tone of their Voice at Unison with it. They first try the Tone of the Glass by striking it, then applying their Mouth near to the Brim of it, sound the same Note with their Voice: this sets the Glass a trembling; they then continually raise their Voice, sounding still the same Note; this encreases the Tremors of the Glass, which by that Means (if it is not too strong) is broke in Pieces.

The

The Effect of *Music* upon Persons bit with the *Tarantula*, (if the Accounts we have of it from Abroad may be credited) is very surprising. A Person bit with the *Tarantula* after some Time loses both Sense and Motion, and dies if destitute of Help. The most effectual Remedy is *Music*. The Musician tries Variety of Airs, till he hits upon one that affects the Patient, who upon that begins to move by Degrees; first keeps Time with his Fingers, Arms, and Legs, afterwards is violently agitated in every Part of his Body; and then leaps up, begins to dance, and increases in Activity every Moment; till after five or six Hours, being very much fatigued, he is put to Bed and left to sweat. The next Day the same Air brings him out of Bed for a new Dance. Which Exercise being thus continued, the Distemper is abated in the Space of four or five Days, the Effects of the Bite being in some Measure carried off by Sweat, and the Patient begins then to recover his Sense and Knowledge by little and little.

The Reason why the Patient is thus affected by the *Music*, is because the Nerves of his Body are so disposed in that Distemper, as easily to be agitated by the Vibrations which are occasioned by the Principle and stronger Notes of what is played.

See on the Subject of this Dissertation, Philosoph. Transact. No. 134, 243, 302, 313, 319, 337. Hist. de l'Acad. 1702, 1708. Grew's Cosmolog. Sacr. Book I. Chap. 5. Mead upon Poisons, p. 59. Keil's Anatomy, p. 214. Baglivi Prax. Medic. Dissert. I. Bellini de Urinis & Pulsibus, &c. sub fine Capitis de Mania.

DISSERTATION II.

Of Capillary Tubes.

BY a *Capillary Tube* is generally understood a Glass Pipe ; the Diameter of whose Bore is at most but about one tenth of an Inch ; though any Tube whose Cavity does not exceed that Magnitude, may be so called.

The Phænomena of Capillary Tubes being such as contradict a known Law in *Hydrostatics*, viz. *that a Fluid rises in a Tube to the same Height with the Level of its Source* * ; and likewise of Affinity with the Ascent of the Sap through the Stems of Plants for the Nourishment of their Fruit, and with divers other Operations of Nature : it has been thought of no small Moment in Philosophy to find out and establish their true Cause ; which after numerous Experiments and several Conjectures about it, is found to be no other than the Attraction of Cohesion ; by which small Particles of Matter mutually adhere together and form larger Bodies†. I shall lay down

* See Chap. I §. 9. Case 4 in the Notes.

† See *Hauksbee's* and *Power's* Experiments. *Musschenbroeck* 4^{to} Edit. *Philosoph Transact.* No. 355 *Mem. de l'Acad.* 1705, 1714, 1722, 1724. With others referr'd to in *Quæstiones Philosoph.*

the several Phænomena, as so many Matters of Fact; and subjoin to each a Solution from that Cause. In Order to which, it may not be improper to premise the following Consideration by Way of *Lemma*.

Let us suppose the Vessel ABCD (*Fig. 15.*) filled with a Fluid to the Height LM, and let it be conceived as divided into the equal Portions EFGH, GHIK, IKLM, &c. farther, let it be supposed, that each Particle of Matter in the inner Surface of the Vessel, has a Sphere of Attraction, whose Semidiameter is equal to the Breadth of three of those Portions; that is to say, that the Attraction of the Particle M reaches upwards as far as F, and downwards as far as S; and that of the Particle O, upwards as far as H, and downwards as far as U; and so of all the rest quite round the Tube. From hence it will follow, that every Particle of the inner Surface of the Vessel, which lies between EF and RS conspires in endeavouring to raise the Fluid towards AB the Top of the Vessel; and that the Fluid is not affected by any other. For Instance, the Particle S, and all below it, will attract downwards three Strata of the Fluid (such as are contained in three equal Divisions of the Vessel) from above, and as many upwards from below; and therefore will have no Effect at all in raising or depressing the Fluid. But the Particle Q will attract only two Strata down-

downwards; because there are no more above it, and three upwards, and therefore will in some Measure tend to raise the Fluid; so the Particle O will attract but one downwards and three upwards; the Particle M none downwards, and three upwards; the Particle K two upwards, and H only one: All which may clearly be seen by their Situations in the Figure with Respect to the Surface of the Fluid. Therefore in every Vessel, where there is a mutual Attraction between the Fluid it contains and the Particles of which it is composed, there will be a certain Number of Particles disposed around it in Form of a broad Periphery or Zone as represented by AB; (*Fig. 16;*) half of which lies above the Surface of the Fluid and half below it, that will tend to make it rise towards the Top. This being understood, the following Phænomena will not be difficult *.

* I have been the more particular in explaining this *Lemma*, because it is not a bare *Periphery* of no Breadth, to which the Ascent of the Fluid is owing, but a *Zone* or *Cingulum* of Particles distended equally in Breadth both ways from the Surface of the Fluid; and because it is upon the Breadth of this, that some of the following Solutions depend. As to the Thickness of it, that undoubtedly is equal to the Semidiameter of the Sphere of Attraction in the Particles of the Vessel; and therefore Vessels, whose Sides are of different Thicknesses (provided those Thicknesses be less than that Semidiameter) must have different Effects upon the same Fluid, though no one has as yet been so accurate as to observe it. The Reason, why a Fluid will not rise in a large Vessel, as well as in one that is Capillary, is because the Attraction of its Particles does not reach far enough into the Middle of the Vessel; and therefore

I. Let there be two Capillary Tubes AB and CD (*Fig. 17.*) open at both Ends; and having their lower Orifices A and C immersed below the Surface of the Water contained in the Vessel FGHI: The Water will immediately rise up in each Tube above the Surface of that in the Vessel, beginning with a swift Motion, which will gradually decrease; till as much Water has entered the Tubes, as they are able to raise: And the Heights to which the Water will rise in them, will be reciprocally as their Diameters.

That the Water ought to rise in both Tubes is an immediate Consequence of the foregoing *Lemma*; because the Column of Water within the Tube is rendered lighter than an equal Column on the outside, as being attracted upwards by a Portion of the interior Surface of the Vessel; and therefore will rise till it becomes as much longer than the external ones as it is made lighter; that the *Æquilibrium*, which was destroyed by the Attraction of the Tube, may be restored by the Weight of the Column. The Reason that the Velocity with which it rises, ought constantly to decrease, is because the heavier the Column is, the less is the Effect of the Attraction, which is always the same in a Tube of the same Diameter.

It only rises about the Sides, standing higher there than in the Middle: As may be seen in a Drinking-Glass, when a Quantity of Water is put into it, somewhat less than is sufficient to fill it.

And

rt II.

s AB
and
mer-
ained
nme-
rface
swift
; till
they
hich
pro-

ubes
oing
with-
qual
up-
e of
be-
nes
m,
the
the
ith
is
is
ys
er.

he
ity
t.
d

Fig. 11. P. 25. Fig. 12. P. 28. Fig. 13. P. 29






Fig. 14. P. 53



Fig. 16. P. 59





Fig. 15. P. 58.



Dis

And
rises
mete
dire
that
Brea
Circ
ame
ing
i. e.

It
in t
than
ther
mer
it st
it v
a pe
ged
qui

T
wh
to t

*
cal E
Tub
Squa
this
a dir
bein
mete
one
Squa

And lastly, the Heights to which the Water rises in them, will be reciprocally as their Diameters; for then the Quantities raised will be directly as the Diameters^{*}; but the Peripheries that raise them, (being always of the same Breadth and having their Lengths equal to the Circumferences of the Tubes,) are as those Diameters; the Quantities of Water therefore, being in the same Ratio, are as the Peripheries, *i. e.* as the Causes by which they are raised.

II. If the Tubes, before they are immersed in the Water, are filled to greater Heights, than those to which it would naturally rise in them; and then have their lower Orifices immersed in Water, the Water will subside till it stands in each at the same Height to which it would have risen; but, if they are held in a perpendicular Position without being immersed, the Water will not subside in the Tubes *quite* so far.

The Reason why the Water in the Tube when its lower Orifice is immersed, subsides to the same Height it would have risen to, had

* The Heights to which the Water rises, being in a reciprocal Ratio of the Diameters; and the Contents of Cylindrical Tubes being in a direct Ratio of their Heights, and of the Squares of their Diameters; the Quantities of Water raised in this Case will be in a reciprocal Ratio of the Diameters, and a direct one of the Squares of the same. Now these two Ratios being compounded together, give the direct one of the Diameters themselves; because the simple reciprocal Ratio destroys one of those, which are contained in the direct one of the Squares.

the Tube been immersed when empty, is because the Column is suspended in one Case by the same Cause, by which it is raised in the other; but when the Tube full of Water is held erect, without being immersed, it will not subside *quite* so far; because the lower End of the Tube, which the Water leaves behind it as it drops out, attracts it the contrary Way; so that the Column in this Case is suspended not only by the inner Surface of the Tube at the Top, but also by its lower End; and therefore a greater Quantity of Water is suspended than in the former Case.

III. If a Tube, having its lower Orifice immersed in Water, be held obliquely, it will raise the Water to the same perpendicular Height, as when held erect.

For since Fluids press according to their perpendicular Heights, the Weight of the Column raised will not be proportionable to the attractive Force of the Tube, till it has arrived at the same perpendicular Height, to which it would have rose, if held erect.

IV. If a Tube, when the Water is risen in to it to its wonted Height, is laid in an Horizontal Situation, the Water will move towards the Middle of the Tube, leaving the End which was immersed a little behind.

The Solution of this Phenomenon depends on what was observed in the Lemma about the Breadth of the attracting Periphery, and its

its being equally situated on each Side the Surface of the Water; for from thence it follows, that if the Water should not run from the full End of the Tube, after it has laid in an Horizontal Situation, but remain contiguous to it; that End of the Column of Water would be attracted only by such a Portion of a Periphery as lies within its Surface at that End; because the End of the Tube coinciding with the Surface, the other Half of the Periphery is wanting. Whereas, at the other End of the Column, there is a Periphery whose Breadth is entire; which overpowering the other, causes the Water to move towards the Middle of the Tube, till as much Periphery is left behind, as can at once affect the Water by the Attraction of its Particles; after which the Water being attracted equally each way, remains at Rest.

V. Let there be a Tube (*Fig. 18.*) consisting of two Parts DR and RCK, of different Diameters; it follows from what has been said, that DR the smaller Part of the Tube, is able to raise Water higher than the other: Let then the Height to which the larger would raise it be TF, and that to which it would rise in the lesser (was it continued down to the Surface of the Fluid) be XL. If this compound Tube be filled with Water and the larger Orifice CK be immersed in the same Fluid the Surface of the Water will sink no farther

farther than XL; the Height to which the lesser Part of the Tube would have raised it.

But if the Tube be inverted as in *Fig. 19.* and the smaller Orifice XL be immersed, the Water will run out till the Surface falls to TF; the Height to which the larger Part of the Tube would have raised it. The Size of the lower Part making no Alteration in the Height, at which the Fluid is suspended in either Case.

In order to account for these Phenomena, it must be considered, that when a Body is so disposed, that its different Parts shall move with different Degrees of Velocity, the greater Proportion the Velocity of that Part to which a moving Power is applied, bears to that of the rest; so much the more effectual is the Power in moving that Body; or that the same Power applied to different Parts, will be equivalent in Effect to different Powers applied to the same Part: As is the known Case of the *Lever*, and all the other *Mechanical* Powers.

Now let us conceive the Tube DR (*Fig. 18.*) continued down to HI, and let it be supposed at present that the Fluids contained in the Tube XLHI and the compound one XLKC, are not suspended by the Periphery at L, but that they press upon their respective Bases HI and CK. Let it farther be supposed that these Bases are each of them moveable
and

and that they are raised up or let down with equal Velocities; then will the Velocity with which XL the uppermost Stratum of the Fluid XLCK moves, exceed that of the same Stratum, considered as the uppermost of the Fluid in the Tube XLHI, as much as the Tube RCK is wider than DR (by the *Lemma* Chap. 1. §. 9.) that is, as much as the Space MNKC exceeds XLIH; consequently by the foregoing Observation, the Effect of the attracting Periphery XL, as it acts upon the Fluid contained in the Vessel XLCK, exceeds its Effect, as it acts upon that in XLHI, in the same Proportion. Since therefore it is able (*ex Hypoth.*) to sustain the Weight of the Fluid XLHI by its natural Power, it is able under this *Mechanical* Advantage, to sustain the Weight of as much, as would fill the Space MNKC: But the Pressure of the Fluid XLCK is equal to that Weight, as having the same Base and an equal Height (Chap. 1. §. 9.) its Pressure therefore, or the Tendency it has to descend in the Tube, is equivalent to the Power of the attracting Periphery XL, for which Reason it ought to be suspended by it.

Again, the Height (*Fig. 19.*) at which the attracting Periphery in the larger Part of the Tube is able to sustain the Fluid is no greater than NF, that to which it would have raised it, had the Tube been continued down to MN. For here the Power of the attracting

ing Periphery acts under a like *Mechanical* Disadvantage; and is thereby diminished in Proportion to the Capacity of the Tube TFXL to that of HIXL; because, if the Bases of these Tubes are supposed to be moved with equal Velocities, the Rise or Fall of the Surface of the Fluid TFXL would be so much less than that of TFMN. And, whereas the attracting Periphery TF is able by its natural Power to suspend the Fluid only to the Height NF in the Tube TFMN; it is in this Case able to sustain no greater Pressure than what is equal to the Weight of the Fluid in the Space HIXL: but the Pressure of the Fluid TFXL which has equal Height and the same Base with it, is equal to that Weight; and therefore is a just *Æquipondium* to the attracting Power.

VI. From hence we may clearly see the Reason, why a small Quantity of Water put into a Capillary Tube, which is of a Conical Form and laid in an Horizontal Situation, will run towards the narrower End. For let AB (*Fig. 20.*) be the Tube, CD a Column of Water contained within it; when the Fluid moves, the Velocity of the End D will be to that of the End C reciprocally as the Cavity of the Tube at D, to that at C (by the *Lem- ma* Chap. 1. §. 9.) that is, reciprocally as the Square of the Diameter at D, to the Square
of

of the Diameter at C^{*}; but the attracting Periphery at D is to that at C, in the simple Ratio of the Diameter at D, to the Diameter at C. Now, since the Effect of the Attraction depends, as much upon the Velocity of that Part of the Fluid where it acts, as upon its natural Force, its Effect at D will be greater than at C; for though the Attraction at D be really less in itself than at C, yet its Loss of Force upon that Account, is more than compensated by the greater Velocity of the Fluid in that Part: the Fluid will therefore move towards B.

VII. From hence likewise it follows, that if a Vessel as ABC (represented *Fig. 21.*) of any Form whatever, have its upper Part drawn out into a Capillary Tube as B; and if this Vessel is filled with Water, and have its lower Orifice placed on FG the Surface of the same Fluid; then the Water will remain suspended in the Vessel, provided the Capillary at the Top be small enough, (was it continued down to the Bottom) to raise the Fluid to the Height B. Because, by Proposition the fifth, the lower Part of the Tube makes no Alteration in the Height, at which the Capillary B is able to sustain the Fluid.

VIII. And if the same Vessel be filled only to the Height DE (*Fig. 22.*) and a Drop of

* 12. *El.* 2.

Water be put into the Capillary at B, (the intermediate Part BDE being full of Air) the Water will continue suspended at the Height DE.

For, although the Fluid ACDE is not in Contact with the Drop of Water in the Capillary Tube, and therefore not immediately supported by it; yet the Pressure of the Atmosphere upon the Surface FG, and against the upper Part of the Drop in the Capillary B keeps the Fluid ADEC, and the Drop and the intermediate Air from separating, just as in the former case the Attraction of Cohesion in the Particles of the Water prevented a Separation between that in the Vessel and that in the Capillary. Consequently, as the Water in the Capillary was able in the former Case to sustain as much Fluid as the Vessel could contain, it is now sufficient to sustain the Fluid ADEC*.

IX. Let there be a Capillary Siphon, as that represented *Fig. 23, 24 or 25*, and let EF be the Height, to which Water might be raised by a Periphery equal to that at A. Now, since

* This happens quite otherwise in Vacuo, because the Pressure of the Air, which as it were connects the Drop with the Water ADEC, being wanting, it immediately falls for Want of a Support. Whereas the former Phenomenon equally succeeds in Vacuo; which shews that the Parts of the Fluid in the Vessel are connected with each other, and with that in the Capillary by their own mutual Attraction of Cohesion, there being nothing else whereby they can be supported.

(as

(as was observed §. 2.) the lower End of a Tube when it is not immersed, causes a longer Column to be suspended than otherwise would be; that is, it supports a short Column besides that which is sustained by the attracting Periphery; let HI be the Height of such a Column as might be suspended by the End C: then, if any of those Tubes are filled with Water, and held as in the Figure (neither End being immersed) the Fluid will run out of the Tube at C, if CD the Difference of the Legs exceeds EF and HI added together, otherwise not.

For the Column AB is a Counterpoise to BD, being of the same perpendicular Height; and therefore it is only the Weight of the Column CD, which determines the Fluid to move; unless that Weight therefore exceeds the Force of the attracting Periphery at A (which the Fluid AB must leave in rising up the Tube) and also what may be supported by the End C, that is, (*ex Hypoth.*) two Columns whose Heights are EF and HI, it cannot run out at C; otherwise it will, as being destitute of a sufficient Support.

X. If the End A is immersed in Water and the other not (supposing the Tube full as before,) it will run out at C, though CD the Difference of the Legs, only exceeds HI. For then the Attraction at A ceases, and there is nothing to support the Column CD, but the

Power the End C has to prevent Drops from falling off it.

XI. Again, if the End C is immersed in Water, (and the other not) it will run out at C, if CD exceeds EF, otherwise not.

For in this Case, there is nothing to support the Column CD, but the attracting Periphery at A, whose Power is supposed able to raise a Column as EF, and no more.

XII. And, if both Ends are immersed (the Tube being supposed full as before) the Water will run out at the lower, which ever it is. For then the Attraction of both Ends ceases; and the longer Column over-balancing the shorter, the Fluid is determined thereby to run out at the lower End.

XIII. If either of the Tubes (*Fig. 23 or 24.*) are small enough to raise the Water from A to B, and if the Orifice A is immersed, the Fluid will rise to B, passing on to C, where it will run out or be suspended according to the foregoing Cases: but if the Periphery at G (*Fig. 25.*) is such as would not support the Fluid higher than AM, it will stop when it comes at G, and only the Part ABG will be filled with it.

For that Fluid which has passed B assists by its Weight the attracting Periphery in raising the Column AB, and therefore it runs down to C. But, if when it comes to G, the Periphery there is not able to support more than AM
the

* He
be made
a Capill
Form, a
will alw
Attractio
the Weig
so much
it be cro

the Difference of the Legs AB and BG, the Fluid must necessarily stop there; since BG is no more than a Counterpoise to MB, and AM is supposed to be as much as the Periphery at G can sustain.

XIV. Tho' a Capillary Tube be shorter than the Height to which its attracting Periphery is able to raise a Fluid, *v. g.* tho' the Tube AB (*Fig. 26.*) be small enough (was it of sufficient Length) to raise Water as high as C; yet when the End A is immersed, the Fluid will not run out at B, but only be suspended at that Height.

For when the Fluid is risen as high as B, it has then no more Periphery above it, to attract it any farther; and if it was forced up a little higher, it would be attracted back again by the End *.

XV. The Ascent of different Fluids in the same Tube is various. *Musschenbroek* has found, that in a Tube in which Water will rise to the Height of twenty six Lines, Oyl of Wormwood will rise but eighteen or nineteen, whereas Urine will rise thirty three or thirty four.

* Hence we see the Absurdity of supposing that a Fluid may be made continually to flow from a lower Place to an higher by a Capillary Tube, as such; for whether the Tube be of such Form, as is represented *Fig. 23, 24, 25, or 26.* The Fluid will always stop when it comes at the higher End; because the Attraction is then in a Direction contrary to its Motion, and the Weight of the whole Fluid, if the Tube is straight, or of so much as is contained in DC the Difference of the Legs, if it be crooked, is likewise an Impediment to it.

The Reason of which is, because some Fluids are attracted more strongly by Glass, than others are. Mercury exhibits Phænomena just the Reverse of the former; for if the End of a Capillary Tube be immersed below the Surface of that Fluid, it will not rise in the Tube to a Level with that on the outside. This is because the Particles of Mercury attract each other more forcibly than they are attracted by those of Glass *.

* See *Jurin's* Dissert. Philosoph. Transl. No. 363.

According to *Musschenbroeck* the Length of the upper Part of a Tube, which is above the Height to which it is able to raise a Fluid, conduces something towards the raising it; and therefore in a longer Tube a Fluid rises higher than in one of the same Dimensions that is shorter; and if a Tube, with so much Fluid contained in it, as it is able to raise, be laid in an Horizontal Situation, the Fluid will run to the Middle of it. But of this I have had no Experience: 'Tis possible that ingenious Professor though very accurate in making Experiments, might herein be deceived. He acknowledges, (Experiment the fifteenth,) that it sometimes happens otherwise.

Other Authors besides those already referred to, that have treated on this Subject, are *Boyle* Exper. Phys. Mech. Exp. 9. *Sturmius* Colleg. Cur. Tentam. 8. *Bernoulli* Gravit. Æth. *Hooke* Microgr. Obser. 6. *Leenuwenhoek* Continuat. Arcan. Nat. Epist. 131. *Sinclair* Art. Gravit.

DISSERTATION III.

Of the Origin of Fountains.

MANY have been the Conjectures of Philosophers concerning the Origin of Fountains; and great Pains have been taken both by the Members of the Royal Society, and those of the Academy of Sciences at *Paris*, in order to ascertain the true Cause of it. It was ARISTOTLE'S Opinion, and held by most of the ancient Philosophers after him, that the Air contained in the Caverns of the Earth, being condensed by Cold near its Surface, was thereby changed into Water; and that it made its Way through, where it could find a Passage. But we have no Experience of any such Transmutation of Air into Water.

Those, who imagine that Fountains owe their Origin to Waters brought from the Sea by subterraneous Ducts, give a tolerable Account how they lose their Saltnefs by Percolation as they pass through the Earth; but find great Difficulty in explaining by what Power the Water rises above the Level of the Sea, to near the Tops of Mountains, where Springs generally abound; it being contrary to

to the Laws of *Hydrostatics*, that a Fluid should rise in a Tube above the Level of its Source. However, they have found two Ways, whereby they endeavour to extricate themselves from this Difficulty. The one is that of DES CARTES, who imagines, that after the Water is become fresh by Percolation, it is raised out of the Caverns of the Earth in Vapour towards its Surface; where meeting with Rocks near the Tops of Mountains in the Form of Arches or Vaults, it sticks to them, and runs down their Sides, (like Water in an Alembic) till it meets with proper Receptacles, from which it supplies the Fountains. Now, this is a mere *Hypothesis* without Foundation or Probability; for in the first Place, we know of no internal Heat of the Earth to cause such an Evaporation; or if that were allowed, yet 'tis quite incredible, that there should be any Caverns so smooth, and void of Protuberances, as to answer the Ends of an Alembic, in collecting and condensing the Vapours together, in every Place where Fountains arise. There are others (as VARENIUS, &c.) who suppose that the Water may rise through the Pores of the Earth, as through Capillary Tubes by Attraction; but hereby they shew, that they are quite unacquainted with what relates to the Motion of a Fluid through such Tubes. For when a Capillary Tube opens in-

to

* See

Differt. 3. *The Origin of Fountains.* 75

to a Cavity at its upper End, or grows larger and larger, so as to cease to be Capillary at that End; the Water will not ascend through that Tube into the Cavity, or beyond where the Tube is Capillary; because there the Force of Attraction is exerted the contrary Way: Nay, if the Cavity is continually supplied with Water, it will be attracted into the Capillary Tube, and run down it, as through a Funnel, if the lower End is immersed in the same Fluid, as in this Case it is supposed to be *.

It has been a generally received Opinion, and much espoused by MARRIOTTE (a diligent Observer of Nature,) that the Rise of Springs is owing to the Rains and melted Snow. According to him, the Rain-Water which falls upon the Hills and Mountains, penetrating the Surface, meets with Clay or Rocks contiguous to each other, along which it runs, without being able to penetrate them, till being got to the Bottom of the Mountain, or to a considerable Distance from the Top, it breaks out of the Ground and forms Springs.

In Order to examine this Opinion, Mr. PERRAULT, DE LA HIRE, and D. SIDELEAU endeavoured to make an Estimate of the Quantity of Rain and Snow, that falls in the Space of a Year, to see whether it would be sufficient to afford a Quantity of Water, equal to

* See the foregoing Dissertation, Sect. 14.

that which is annually discharged into the Sea by the Rivers. The Result of whose Inquiries was, that the Quantity of Rain and Snow which fell in a Year into a Cylindrical Vessel, would fill it (if secured from evaporating) to the Height of about nineteen Inches. Which Quantity D. SIDELEAU * shewed, was not sufficient to supply the Rivers; for that those of *England, Ireland, and Spain* discharge a greater Quantity of Water annually, than the Rain, according to that Experiment, is able to supply. Besides which, another Observation was made by them at the same Time, viz. that the Quantity of Water raised in Vapour one Year with another, amounted to about Thirty two Inches, which is thirteen more than falls in Rain: A plain Indication, that the Water of Fountains is not supplied by Rains and melted Snow.

Thus, the true Cause of the Origin of Fountains remained undiscovered, till Dr. HALLEY in making his Celestial Observations upon the Tops of the Mountains at *St. Helena*, about eight Hundred Yards above the Level of the Sea, found that the Quantity of Vapour which fell there (even when the Sky was clear) was so great that it very much impeded his Observations, by covering his Glasses with Water every half Quarter of an Hour; and upon that

* Memoirs of the Royal Academy of Sciences for the Year 1632.

Differt. 3. *The Origin of Fountains.* 77

attempted to determine by Experiment the Quantity of Vapour, exhaled from the Surface of the Sea, as far as it arises from Heat; in order to try, whether that might be a sufficient Supply for the Water continually discharged by Fountains. The Process of his Experiment was as follows. He took a Vessel of Water salted to the same Degree with that of Sea-Water, in which he placed a Thermometer; and by means of a Pan of Coals, brought the Water to the same Degree of Heat, which is observed to be that of the Air in our hottest Summer: This done, he affixed the Vessel of Water with the Thermometer in it, to one End of a Pair of Scales, and exactly counterpoised it with Weights on the other. Then, at the End of two Hours, he found by the Alteration made in the Weight of the Vessel, that about a sixtieth Part of an Inch of the Depth of the Water, was gone off in Vapour; and therefore in twelve Hours, one tenth of an Inch would have gone off. Now this accurate Observer allows the *Mediterranean Sea* to be forty Degrees long and four broad; (the broader Parts compensating for the narrower;) so that its whole Surface is one Hundred and sixty square Degrees; which according to the Experiment must yield at least five Thousand two Hundred and eighty Millions of Tons. In which Account, no Regard is had to the Wind, and the Agitation of the

Surface of the Sea; both which undoubtedly promote the Evaporation.

It remained now, to compare this Quantity of Water, with that, which is daily conveyed into the same Sea by the Rivers. The only Way to do which, was to compare them with some known River; and accordingly he takes his Computation from the River *Thames*, and to avoid all Objection, makes Allowances, probably greater than what were absolutely necessary.

The *Mediterranean* receives the following considerable Rivers, viz. the *Iberus*, the *Rhone*, the *Tibur*, the *Po*, the *Danube*, the *Neister*, the *Borystenes*, the *Tanais*, and the *Nile*. Each of these he supposes to bring down ten Times as much Water as the *Thames*; whereby he allows for smaller Rivers, which fall into the same Sea. The *Thames* then he finds by Mensuration to discharge about 20,300,000 Tons of Water a Day. If therefore the abovesaid nine Rivers yield ten Times as much Water as the *Thames* doth, it will follow, that all of them together yield but 1827 Millions of Tons in a Day; which is but little more than one Third of what is proved to be raised in Vapour out of the *Mediterranean* in the same Time. We have therefore from hence a Source abundantly sufficient for the Supply of Fountains.

Now having found, that the Vapour exhaled from the Sea, is a sufficient Supply for the

Foun-

Differt. 3. *The Origin of Fountains.* 79

Fountains; he proceeds in the next Place to consider the Manner in which they are raised; and how they are condensed into Water again; and conveyed to the Sources of Springs.

In Order to this he considers, that, if an Atom of Water was expanded into a Shell or Bubble, so as to be ten Times as big in Diameter as when it was Water, that Atom would become specifically lighter than Air; and therefore would rise so long as the Warmth which first separated it from the Surface of the Water should continue to distend it to the same Degree; and consequently, that Vapours may be raised from the Surface of the Sea in that Manner, till they arrive at a certain Height in the Atmosphere, at which they find the Air of equal specific Gravity with themselves. Here they will float, till being condensed by Cold, they become specifically heavier than the Air, and fall down in Dew, or being driven by the Winds against the Sides of the Mountains, (many of which far surpass the usual Height to which the Vapours would of themselves ascend) are compelled by the Stream of the Air to mount up with it to the Tops of them: where being condensed into Water they presently precipitate, and gleeing down by the Crannies of the Stone, Part of them enters into the Caverns of the Hills; which being once filled, all the over-plus

plus of Water that comes thither, runs over by the lowest Place, and breaking out by the Sides of the Hills, forms single Springs. Many of these running down by the Valleys between the Ridges of the Hills, and coming to unite, form little Rivulets or Brooks: Many of these again meeting in one common Valley, and gaining the plain Ground, being grown less rapid, become a River; and many of these being united in one common Channel, make such Streams as the *Rhine* and the *Danube*; which latter, he observes, one would hardly think to be a Collection of Water condensed out of Vapour, unless we consider how vast a Tract of Ground that River drains, and that it is the Sum of all those Springs, which break out on the South Side of the *Carpathian* Mountains, and on the North Side of the immense Ridge of the *Alps*; which is one continued Chain of Mountains from *Switzerland* to the *Black-Sea*.

Thus one Part of the Vapours, which are blown on the Land, is returned by the Rivers into the Sea, from whence it came; another Part falls into the Sea before it reaches the Land; and this is the Reason, why the Rivers do not return so much Water into the *Mediterranean* as is raised in Vapour. A third Part falls on the Low-Lands, and is the *Pabulum* of Plants, where yet it does not rest; but is again exhaled by Vapour by the
Action

* T
alterna
thence
ciproca
Let
Cavity
as this
lating t
whatev
vity, ti
Duct;
other P
KI it w
our at A
vided th

Differt. 3. *The Origin of Fountains.* 81

Action of the Sun, and is either carried by the Winds to the Sea, to fall in Rain or Dew there, or else to the Mountains to become the Sources of Springs.

However, it is not to be supposed, that all Fountains are owing to one and the same Cause; but that some proceed from Rain and melted Snow, which subsiding through the Surface of the Earth, makes its Way into certain Cavities and thence issues out in the Form of Springs; because the Waters of several are found to increase and diminish in Proportion to the Rain which falls: That others again, especially such as are salt, and spring near the Sea-Shore, owe their Origin to Sea-Water percolated through the Earth; and some to both these Causes: Though without doubt most of them, and especially such as spring near the Tops of high Mountains, receive their Waters from Vapours, as before explained *.

* There is a certain Species of Springs which ebb and flow alternately, and some that cease to flow for a Time, which from thence are called reciprocating or intermitting ones. Their Reciprocations may be accounted for in the following Manner.

Let ABC represent one Side of an Hill in which there is a Cavity DEF, and from this a subterraneous Duët IKL. Now as this Cavity fills with Water (suppose from Vapours percolating through the Surface of the Hill, or in any other Manner whatever) its Surface will rise in the Duët as it does in the Cavity, till it arrives at M, the Level with the upper Part of the Duët; at which Time it will run over at K, filling KLA the other Part of the Duët. Now, if the Column KL is longer than KI it will overpoise the other, and so cause the Water to run out at A, till its Surface in the Cavity sinks as far as I, (provided the Duët is large enough to convey the Water away faster than

than it enters the Cavity) at which Time the Fountain at A will cease to play, till the Surface of the Water in the Cavity rises again to M, and runs over at K as before. The Reason why the Water continues running (when the Duct is once full) till its Surface sinks to I, is because the Air pressing against it as it runs out at A and also upon its Surface in the Cavity, keeps the Duct full, as long as the Water in the Cavity is high enough to feed its Orifice at I.

See more on this Subject in *Philosoph. Transact.* No. 119, 189, 192, 384, 424. *History de l' Acad.* 1693, 1703, 1713. *Gulielmini della Natura de Fiumi.* *Dale's History of Harwich.* *Marriotte's Hydrostatics.* *Nieuwentyt Contempt.* 19. *Varenus Geograph.* Cap. 16. *Regnault Vol. 2. Conversat.* 6. *Hales's Statical Essays Vol. 1. Experiment* 19. *Michelottus in Append ad J. Bernoullii de Effervesc.*



l.

A
ty
on
ll)
it
ps
gh

r 9,
13.
ich.
nius
les's
nd

I S-

Fig. 17.
P. 60.




Fig. 18.
P. 63.



Fig. 19.
P. 64.




Fig. 20. *P. 66.*



Fig. 21. *P. 67.*




Fig. 22. *P. 67.*




Fig. 23.




Fig. 24.
P. 68.



Fig. 25. *P. 71.*



Fig. 26.





Fig. 27.
P. 81.





C

C

I.

II.

III.

IV.

S

V.

Re

F

Prin

A
COMPENDIOUS SYSTEM
OF
Natural Philosophy.
With NOTES,

Containing the MATHEMATICAL
DEMONSTRATIONS, and some
occasional REMARKS.

PART II. Continued.

Consisting of five *DISSERTATIONS*.

- I. Of the *Barometer*.
 - II. Of the Cause and Origin of the *Winds*.
 - III. Of the Ascent of *Vapours*, and their Resolution into
Rain, Hail, Snow, &c.
 - IV. Of the Causes of *Thunder* and *Lightning*, with a
Solution of the *Phænomena* of the *Aurora Borealis*.
 - V. A New Theory of *Fermentation*.
-

By J. ROWNING, M. A.

Rector of ANDERBY in LINCOLNSHIRE, and late
Fellow of MAGDALEN College in CAMBRIDGE.

The SECOND EDITION.

L O N D O N :

Printed for SAM. HARDING, Bookseller, on the
Pavement in St. Martin's Lane.

MDCCLXXXVIII.

Diff

I

IN
tice
rom
Sufp
to t
to
gina
diff
finc



*
to th
inacc
may
than
the C
tion
prop
supp
the C
some
Hea
sequ
over
infe
ever
pole
fore
was

DISSERTATION IV.

Of the Barometer.

IN treating of the Properties of the Air (Chap. III.) I have already taken Notice of the Construction of the common Barometer; and proved, that the Ascent and Suspension of the Mercury therein, is owing to the Pressure * of the Air. I proceed now to a more particular Inquiry into the Original, and Use of this Instrument; and the different Forms under which it has appeared, since the Time of its Inventor *Torricelli*.

In

* To say the Ascent and Suspension of the Mercury is owing to the Pressure and Elasticity of the Air, as is commonly done, is inaccurate. The Variation, indeed, in the Height of the Mercury, may be ascribed to the Elasticity of the Air, but no otherwise, than as to its *remote* Cause; viz. as it occasions an Alteration in the Quantity of Air, impending over the Place where the Variation happens; which alters its Weight, and so the Mercury is proportionably raised or depressed. To illustrate this, let it be supposed, that the Air is every where in *Æquilibrio*, quite round the Globe, and at perfect Rest; and then, that its Elasticity, in some one Place near the Surface of the Earth, is augmented by the Heat of the Sun, all the rest of it remaining as before. The Consequence of this will be, that the superior Part of the Atmosphere, over this Place, will be raised higher by the Expansion of the inferior Air; and therefore, being unconfined, will spread itself, every way, over the neighbouring Columns, which we suppose to retain their former State. The Quantity of Matter therefore in those Columns of Air, in whose lower Parts its Elasticity was increased, will be diminished, and that of the neighbouring

In the Beginning of the last Century, it was a prevailing Opinion among Philosophers, that the Universe was full of Matter; and that Nature (as they expressed it) abhorred a *Vacuum*: Accordingly they imagined, that if a Fluid was sucked up a Pipe with a sufficient Force, it would rise to any Height whatever; since Nature would not suffer any Part of the Pipe to remain empty. *Galilæo*, who flourished about that Time, found upon Trial, that the common Pump would not raise Water, unless the Sucker reached within three and thirty Feet of its Surface in the Well *: From hence

ones augmented. A Barometer therefore placed in those Regions, where the Air was rarified, will subside; while one in the neighbouring Countries will ascend; and they will continue at different Heights, till the denser Air, rushing in upon the rarified, restores the *Æquilibrium*. Thus, we see, the Variation of the Air's Elasticity is not the *immediate* Cause of the Variation in the Barometer; it first affects the Weight of the Air, by altering the Quantity incumbent over any Place, and that affects the Barometer. But, if we may have Recourse to *remote* Causes, we may, if we please, go one Step farther; and say, the Ascent and Suspension of the Mercury is owing to the Heat of the Sun; for by the foregoing Instance, a Variation in the Heat of the Sun may sometimes be the occasion of a Variation in the Height of the Mercury.

Neither is the Suspension of the Mercury, in a Tube, that is kept within Doors, to be ascribed to the Elasticity of the Air; for that exerts no Force, but as the internal Air is pressed by the external, which endeavours to get in, where-ever it can find a Way.

* It is a common Notion, that a sucking Pump will not raise Water above thirty-three Feet, whereas it will raise it to any Height whatever, if the Sucker reaches within thirty-three Feet of the Surface of the Water; as will be evident to any one that considers the Structure of the Pump: For all the Water, which has

once

hence he judiciously inferred, that a Column of Water thirty-three Feet high was a Counterpoise to a Column of Air of an equal Base, whose Height extended to the Top of the Atmosphere; and that, for this Reason, the Water would not follow the Sucker any farther. *Torricelli*, observing this, took the Hint; and considered, that, if a Column of Water, of about thirty-three Feet, was equal in Weight to a Column of Air, of the same Base *; a Co-

once passed through the Valve in the Sucker, is supported by that, as the Sucker is drawn up, and rests upon a Valve placed in the Pump below, as it is let down; so that it can be no Impediment to the rising of the Water below the Sucker, whatever the Length of the Column, which it forms, may be. The placing one Pump above another, where Water is to be raised from great Depths, is rather for Strength and Conveniency, than out of Necessity.

* Perhaps it may be enquired here, how it comes to pass, that the Column of Air, which presses upon the stagnant Mercury in the Bason, is always supposed to have an equal Base with the suspended Column in the Tube; whereas, in Reality, its Base is equal to the Surface of the stagnant Mercury. The Reason is, that, as the Base of the Column of Air increases, in the same Proportion the Velocity, wherewith it descends, decreases, when it forces down the Surface of the Mercury in the Bason; consequently its Moment, or Pressure upon the Surface of the stagnant Mercury (so far as it relates to the suspending of it in the Tube) is no greater, than it would have been, had its Base been equal to that of the suspended Column; and therefore, in considering it as suspending a Fluid in a Tube, it is properly enough said to be a Column of such a Base.

Neither is this Supposition inconsistent with the ninth Proposition of the first Chapter, where it is demonstrated, that the Pressure of a Fluid is in Proportion to its perpendicular Height, and the Quantity of Surface, against which it presses. For, as the Surface of the Mercury may be considered as a Base on which the

Column of Mercury, no longer than about twenty-nine Inches and a half, would be so too; such a Column of Mercury being as heavy, as thirty-three Feet of Water. Accordingly he tried the Experiment in a Glass Tube (in the Manner laid down, Chap. III. §. 4.) and found it to succeed *. The Apparatus he made Use of,

Column of Air rests, so the Base of the Column of Air may be consider'd as a Surface against which the Mercury presses. These two being equal, 'tis clear, that only the Relation of the Heights of the Columns are to be considered, and not that of their Bases.

* Notwithstanding this clear Proof of the Pressure of the Atmosphere, the Asserters of a *Plenum* would by no Means be prevailed upon to allow it to be such; but tried all Ways to account for this *Phænomenon* from some other Cause. The most chimerical Solution, and which at the same Time gave the adverse Party the greatest Difficulty to overthrow, was that of *Linus*. He contended, that in the upper Part of the Tube, there is a Film, or Rope of Mercury, extended thro' the seeming Vacuity, and that the rest was suspended by it, and kept from falling into the Bason; and that this Film is able to support about twenty-nine Inches of Mercury. He confirms his *Hypothesis* by the following Experiment. Take, says he, a small Tube, open at both Ends, suppose about twenty Inches long; fill this Tube with Mercury, stopping the lower Orifice with your Thumb: Then closing the upper with your Finger, and immersing the lower in stagnant Mercury, you shall perceive, upon the Removal of your Thumb, a manifest Suction of your Finger into the Tube; and the Tube and Mercury will both stick so close to it, that you may carry them about the Room. Therefore, says he, the internal Cylinder of Mercury in the Tube is not held up by the preponderant Air without; for if so, whence comes so strong a Suction, and so firm an Adhesion of the Tube to your Finger?

Or if you fill the same Tube almost full of Mercury, leaving a little Space of Air within, and then immerse it in the stagnant Mercury, you will find, that, notwithstanding its Surface is at some Distance from your Finger, there will be a considerable Suction,

Differt. IV. Of the Barometer. 87

of, is now the common Barometer or Weather Glas *.

The Mercury standing at a less Height, the nearer it is carried to the Top of the Atmosphere,

tion of it, as before. From hence he infers, that the Finger supports the Mercury, by Means of the abovementioned Film, and that the Pressure of the Atmosphere is not concerned.

But, when it was found, that the Mercury would not stand so high in the Tube, on the Top of a Mountain, as below; and would quite fall, when the circumambient Air was extracted from it by the Pump, all Objections vanished; and *Linus's* funicular Hypothesis (as it was called) though it seemed to solve all other Phenomena, relating to the Suspension of the Mercury, was with Justice rejected.

Kircher, when this new Doctrine of a Vacuum was first advanced at Rome, contended, that the Authors of it were establishing Principles not only repugnant to those of Nature, but such as would be prejudicial to the Orthodox Faith; as endeavouring to evince by this subtle Experiment, that there might be in Nature *beatum sine loco, accidentia sine subjecto*, and therefore made the Experiment with Water, in the following Manner. He caused a small Bell to be fixed in the upper Part of the Tube, imagining, that, if there should be a Vacuum, the Bell would not be made to sound: But in making the Experiment, some Air got into the Tube (for he tells us, that but ten Feet of Water remained in the Tube, after it was inverted) the Bell therefore was heard to sound; and so the Notion of a Vacuum, till more accurate Experiments evinced the contrary, was exploded with Contempt.

* *Huygens* observed, that, if a Tube seventy-five Inches long, was filled with Mercury well purged of its Air, the whole Quantity of Mercury would remain suspended; whereas, according to the *Torricellian* Experiment, the Mercury ought to have subsided to the Height of about twenty-nine Inches.

The Cause of this Phenomenon seems to be, that, by the great Weight of so long a Column of Mercury, it was pressed into so close Contact with the Glas in pouring in, that by the mutual Attraction of Cohesion between the Mercury and the Glas, the whole Column was sustained, after the Tube was inverted.

(Chap.

(Chap. III. §. 7.) renders it useful in determining the Height of Mountains; and finding out the different Elevation of one Place above another. Accordingly, Dr. *Halley* has given us a Table for that Purpose, in the Philosophical Transactions N^o. 181, shewing how many Feet the Descent of the Mercury each Inch answers to, as it is conveyed to the Top of a Mountain, or other elevated Place. And Dr. *Nettleton* has done the like in the Philosophical Transactions N^o. 388, shewing what Number of Feet answers to each tenth Part of an Inch, from twenty-six to thirty-one Inches of Mercury.

But the principal Use of it is, to estimate the Gravity of the Air at different Times, in Order to foresee the Alterations of the Weather, which are consequent thereon. To this End, Dr. *Halley* in the same Transaction has also laid down the more remarkable *Phænomena*, relating to the different Heights of the Mercury at different Times, together with the Solution of each; which are so just, and so agreeable to true Philosophy, that I doubt not but the Reader will excuse me for giving his Account in his own Words, rather than to render it imperfect, by endeavouring to vary from it, or abridge it.

“ 1. In calm Weather, when the Air is inclined to Rain, the Mercury is commonly low.

“ In

Differt. IV. *Of the Barometer.* 89

“ 2. In serene, good, settled Weather, the
“ Mercury is generally high.

“ 3. Upon very great Winds, though they
“ be not accompanied with Rain, the Mercury
“ sinks lowest of all, with Relation to the
“ Point of the Compass the Wind blows upon.

“ 4. *Cæteris paribus*, the greatest Heights
“ of the Mercury are found upon easterly and
“ north-easterly Winds.

“ 5. In calm frosty Weather, the Mercury
“ generally stands high.

“ 6. After very great Storms of Wind,
“ when the Mercury has been low, it gene-
“ rally rises again very fast.

“ 7. The more northerly Places have greater
“ Alterations of the Barometer, than the more
“ southerly.

“ 8. Within the Tropics, and near them,
“ those Accounts we have had from others,
“ and my own Observations at *St. Helena*,
“ make very little or no Variation of the
“ Height of the Mercury in all Weathers.

“ Hence I conceive that the principal Cause
“ of the Rise and Fall of the Mercury, is from
“ the variable Winds, which are found in the
“ temperate Zone, and whose great Uncon-
“ stancy, here in *England*, is most notorious.

“ A second Cause is the uncertain Exhalation
“ and Precipitation of the Vapours lodging in
“ the Air, whereby it comes to be at one Time,
“ much more crouded than at another, and

“ con-

“ consequently heavier, but this latter in a great
“ Measure depends upon the former. Now,
“ from these Principles, I shall endeavour to
“ explicate the several *Phænomena* of the Ba-
“ rometer, taking them in the same Order I
“ laid them down. Thus.

“ 1. The Mercury's being low, inclines it
“ to rain, because the Air being light, the
“ Vapours are no longer supported thereby,
“ being become specifically heavier than the
“ Medium wherein they floated, so that they
“ descend towards the Earth, and in their Fall,
“ meeting with other aqueous Particles, they
“ incorporate together, and form little Drops
“ of Rain; but the Mercury's being at one
“ Time lower than at another, is the Effect of
“ two contrary Winds blowing from the Place
“ where the Barometer stands; whereby the
“ Air of that Place is carry'd both Ways from
“ it, and, consequently, the incumbent Cylin-
“ der of Air is diminished, and accordingly the
“ Mercury sinks: As for Instance, if in the
“ *German* Ocean it should blow a Gale of
“ westerly Wind, and at the same Time an
“ easterly Wind in the *Irish* Sea; or if in
“ *France* it should blow a northerly Wind,
“ and in *Scotland* a southerly; it must be
“ granted, that that Part of the Atmosphere
“ impendant over *England*, would thereby be
“ exhausted and attenuated, and the Mercury
“ would subside, and the Vapours, which be-
“ fore

“ fore floated in those Parts of the Air of
 “ equal Gravity with themselves, would sink
 “ to the Earth.

“ 2. The greater Height of the Barometer
 “ is occasioned by two contrary Winds blow-
 “ ing towards the Place of Observation, where-
 “ by the Air of other Places is brought thither
 “ and accumulated; so that the incumbent
 “ Cylinder of Air, being encreased both in
 “ Height and Weight, the Mercury pressed
 “ thereby must needs stand high, as long as
 “ the Winds continue so to blow; and then
 “ the Air being specifically heavier, the Va-
 “ pours are better kept suspended, so that they
 “ have no Inclination to precipitate and fall
 “ down in Drops, which is the Reason of the
 “ serene good Weather which attends the
 “ greater Heights of the Mercury.

“ 3. The Mercury sinks the lowest of all by
 “ the very rapid Motion of the Air in Storms
 “ of Wind. For the Tract or Region of the
 “ Earth’s Surface, wherein the Winds rage,
 “ not extending all round the Globe, that
 “ stagnant Air which is left behind, as like-
 “ wise that on the Sides, cannot come in so
 “ fast as to supply the Evacuation made by so
 “ swift a Current, so that the Air must neces-
 “ sarily be attenuated, when and where the
 “ said Winds continue to blow, and that more
 “ or less, according to their Violence; add
 “ to which, that the horizontal Motion of the

“ Air being so quick as it is, may, in all Pro-
 “ bability, take off some Part of the perpendi-
 “ cular Pressure thereof; and the great Agi-
 “ tation of its Particles is the Reason why the
 “ Vapours are dissipated, and do not condense
 “ into Drops, so as to form Rain, otherwise
 “ the natural Consequence of the Air’s Rare-
 “ faction †.

“ 4. The Mercury stands the highest upon
 “ the easterly and north-easterly Wind, be-
 “ cause in the great *Atlantic* Ocean, on this
 “ Side the thirty-fifth Degree of north Lati-
 “ tude, the Winds are almost always wester-
 “ ly or south-westerly; so that whenever
 “ here the Winds come up at east and north-
 “ east, ’tis sure to be checked by a contrary
 “ Gale as soon as it reaches the Ocean; where-
 “ fore, according to what is made out in our
 “ second Remark, the Air must needs be heap-

† The Reason the Doctor assigns for the sinking of the Mercury the lowest of all in violent Storms of Wind, seems not sufficient. Perhaps it may be better accounted for thus; the Cause why the Wind blows at all, is in order to restore the *Æquilibrium* of the Atmosphere, when lost (as may be inferred from what was said in the first Note of this, and will be more largely explained in the following Dissertation;) it therefore always blows towards that Point, where the Air is most rarefied and lightest. Now the Air in its Progress to that Point, must certainly move faster and faster; for the Cause which gave it Motion at first, continues to act upon it all the Way. Consequently, in whatever Place the Wind blows with great Rapidity, that Place is at, or near the Point, where the Air is most rarefied, and lightest; which is a sufficient Reason for the Mercury’s standing low at that Place.

Differt. IV. *Of the Barometer.* 93

“ ed over this Island, and consequently the
 “ Mercury must stand high, as often as these
 “ Winds blow. This holds true in this Coun-
 “ try, but is not a general Rule for others,
 “ where the Winds are under different Cir-
 “ cumstances; and I have sometimes seen the
 “ Mercury here, as low as twenty-nine Inches
 “ upon an easterly Wind, but then it blew ex-
 “ ceeding hard, and so comes to be accounted
 “ for, by what was observed upon the third
 “ Remark.

“ 5. In calm frosty Weather the Mercury
 “ generally stands high, because (as I con-
 “ ceive) it seldom freezes, but when the Winds
 “ come out of the northern, and north-eastern
 “ Quarters; or, at least, unless those Winds
 “ blow at no great Distance off: For the
 “ north Parts of *Germany, Denmark, Swe-*
 “ *den, Norway,* and all that Tract from
 “ whence north-eastern Winds come, are
 “ subject to almost continual Frost all the
 “ Winter; and thereby the lower Air is very
 “ much condensed, and in that State is brought
 “ thitherwards by those Winds, and being ac-
 “ cumulated by the Opposition of the wester-
 “ ly Wind blowing in the *Ocean*, the Mer-
 “ cury must needs be pressed to a more than
 “ ordinary Height: and, as a concurring
 “ Cause, the shrinking of the lower Parts of
 “ the Air into lesser Room by Cold, must
 “ needs

“ needs cause a Descent of the upper Parts of
 “ the Atmosphere, to reduce the Cavity made
 “ by this Contraction to an *Æquilibrium*.

“ 6. After great Storms, when the Mer-
 “ cury has been very low, it generally rises
 “ again very fast: I once observed it to rise
 “ one Inch and a half in less than six Hours,
 “ after a long continued Storm of south-west
 “ Wind. The Reason is, because the Air be-
 “ ing very much rarefied, by the great Eva-
 “ cuations which such continued Storms make
 “ thereof, the neighbouring Air runs in the
 “ more swiftly, to bring it to an *Æquilibri-*
 “ *um*; as we see Water runs the faster for
 “ having a greater Declivity.

“ 7. The Variations are greater in the more
 “ northerly Places, as at *Stockholm*, greater
 “ than that at *Paris* (compar'd by M. *Paf-*
 “ *chal*;) because the more northerly Parts
 “ have usually greater Storms of Wind than
 “ the more southerly, whereby the Mercury
 “ should sink lower in that Extream; and
 “ then the northerly Winds bringing the more
 “ dense and ponderous Air from the Neigh-
 “ bourhood of the Pole, and that again being
 “ checked by a southerly Wind at no great
 “ Distance, and so heaped, must of Necessity
 “ make the Mercury in such Case stand higher
 “ in the other Extream.

“ 8. Lastly,

“ 8. Lastly, this Remark, That there is
 “ little or no Variation near the *Equinoctial*,
 “ does above all others, confirm the *Hypothese*
 “ of the variable Winds being the Cause of
 “ these Variations of the Height of the Mer-
 “ cury ; for in the Places above-named, there
 “ is always an easy Gale of Wind blowing
 “ nearly upon the same Point, *viz.* east-north-
 “ east, at *Barbadoes*, and east-south-east at
 “ *St. Helena* *, so that there being no contra-
 “ ry Currents of Air to exhaust or accumulate
 “ it, the Atmosphere continues much in the
 “ same State : However, upon Hurricanes
 “ (the most violent of Storms) the Mercury
 “ has been observed very low, but this is but
 “ once in two or three Years, and it soon re-
 “ covers its settled State about $29 \frac{1}{2}$ Inches.”

Monfieur *Leibnitz* accounted for the De-
 scent of the Mercury before Rain, upon an-
 other Principle †, *viz.* as a Body specifically
 lighter than a Fluid, while it is suspended by
 it, adds more Weight to that Fluid, than
 when, by being reduced in its Bulk, it be-
 comes specifically heavier, and descends ; so
 the Vapour, after it is reduced into the Form
 of Clouds, and descends, adds less Weight to
 the Air, than before ; and therefore the Mer-

* See the Cause of this assigned in the following Dissertation.

† *Memoir. de l'Acad.* 1711.

cury falls. To which it is answered, 1st. That when a Body descends in a Fluid, its Motion, in a very little Time, becomes uniform, (or nearly so) a farther Acceleration of it being prevented by the Resistance of the Fluid; and then, by the third Law of Nature, it presses the Fluid downwards, with a Force equal to that whereby it tends to be farther accelerated, that is, with a Force equal to its whole Weight. 2^{dly}. The Mercury, by its Descent, foretells Rain a much longer Time before it comes, than the Vapour, after it is condensed into Clouds, can be supposed to take up in falling. 3^{dly}. Supposing that as many Vapours, as fall in Rain, during the Space of a whole Year, were at once to be condensed into Clouds, and even quite cease to gravitate upon the Air, its Gravity would scarce be diminished thereby, so much as is equivalent to the Descent of two Inches of Mercury in the Barometer. Farther, in many Places between the Tropics, the Rains fall at certain Seasons, in very great Quantities *, and yet the Barometer shews there very little or no Alteration in the Weight of the Air.

The following are Mr. *Patrick's* Observations on the rising and falling of the Mercury. They are very just, and are to be accounted

* See Dissertation the vith.

Differt. IV. *Of the Barometer.*

97

for on the same Principles with those of Dr. Halley.

“ 1. The rising of the Mercury presages in
“ general fair Weather ; and its falling, foul
“ Weather ; as Rain, Snow, high Winds and
“ Storms.

“ 2. In very hot Weather, the falling of
“ the Mercury foreshews Thunder.

“ 3. In Winter the rising presages Frost ;
“ and in frosty Weather, if the Mercury falls
“ three or four Divisions, there will *certain-*
“ *ly* follow a Thaw. But in a continued
“ Frost, if the Mercury rises, it will *certain-*
“ *ly* snow.

“ 4. When foul Weather happens soon after
“ the Falling of the Mercury, expect but little
“ of it. And, on the contrary, expect but
“ little fair Weather, when it proves fair
“ shortly after the Mercury has risen.

“ 5. In foul Weather, when the Mercury
“ rises much and high, and so continues for
“ two or three Days before the foul Weather
“ is quite over, then expect a Continuance of
“ fair Weather to follow.

“ 6. In fair Weather, when the Mercury
“ falls much and low, and thus continues for
“ two or three Days before the Rain comes ;
“ then expect a great deal of wet, and pro-
“ bably high Winds.

“ 7. The

“ 7. The unsettled Motion of the Mercury denotes uncertain and changeable Weather.

“ 8. You are not so strictly to observe the Words engraven on the Plates (though, for the most Part, it will agree with them) as the Mercury’s *Rising* and *Falling*: For if it stands at *Much Rain*, and then rises up to *Changeable*, it presages fair Weather, although not to continue so long, as it would have done, if the Mercury were higher: And so on the contrary, if the Mercury stood at *Fair*, and falls to *Changeable*, it presages foul Weather; though not so much of it, as if it had sunk down lower.”

From these Observations, it appears, That it is not so much the Height of the Mercury in the Tube, that indicates the Weather, as the Motion of it up and down; wherefore, in Order to pass a right Judgment of what Weather is to be expected, we ought to know, whether the Mercury is actually *Rising* or *Falling*, to which End, the following Rules are of Use.

1. If the Surface of the Mercury is convex, standing higher in the Middle of the Tube than at the Sides, it is generally a Sign that the Mercury is then rising.

2. If the Surface is concave, or hollow in the Middle, it is sinking. And,

3. If

Differt. IV. *Of the Barometer.* 99

3. If it is plain, the Mercury is stationary, or rather, if it is a little convex; for Mercury being put into a Glass Tube, especially a small one, will naturally have its Surface a little convex; because the Particles of Mercury attract each other more forcibly than they are attracted by Glass. Further,

4. If the Glass is small, shake the Tube; and if the Air is growing heavier, the Mercury will rise about half the tenth of an Inch higher, than it stood before; if it is growing lighter, it will sink as much. This proceeds from the Mercury's sticking to the Sides of the Tube, which prevents the free Motion of it, till it is disengaged by the Shock. And therefore, when an Observation is to be made with such a Tube, it ought always to be shaken first, for sometimes the Mercury will not vary of its own Accord, till the Weather, it ought to have indicated, is present.

The Usefulness of knowing, whether the Mercury is actually rising or falling; and the Advantage that would arise from perceiving the most minute Variations in estimating the Heights of Places, have given Occasion to the Invention of several Kinds of Barometers different from the *Torricellian*, though founded on the same Principle; wherein the Scale of Variation, which in that is not above three Inches,

O

should

should be considerably larger. Of which I am now to give some Account.

1. The first is that of *Des Cartes*, which was made in the Form expressed *Fig. 28.* where AB is a Tube hermetically * sealed at A, and having its lower Orifice immersed in stagnant Mercury EF, and filled with the same Fluid to G, from thence to H with Water, and empty from thence to the Top. Now, when the Mercury rises in this Tube, suppose from G to L, the Water will be raised in the small Tube, perhaps from H to M, *viz.* as many Times further, as the Tube CA is smaller than CD; by which Means the Variations become much more sensible, than they are in the common Barometer. The Inconvenience of this was, that the Air, included in the Water, getting loose by Degrees, filled the void Space at the Top, and so spoiled the Machine.

2. He then contrived it thus, ABC (*Fig. 29.*) is a bent Tube hermetically sealed at A, filled with Water from F to D (tinged with *Aqua Regia* to prevent its freezing,) from D to E with Mercury, and empty from thence to the Top. Then, upon the Mercury's rising, suppose from

* A Tube is said to be *hermetically* sealed, when the End is so closed, that nothing can possibly evaporate through it. And, because this is best done, when it is closed up with its own Substance; or when its Bore does not reach quite through it, it is then said to be *hermetically* sealed.

Differt. IV. *Of the Barometer.* 101

E to M, and falling as much at D, the Surface of the Water at F would sink so many Times farther than the Surface of the Mercury at D, as the Tube CG was smaller than GH. The Water here is liable to evaporate, though that may, in some Measure, be prevented, by pouring a few Drops of Oil of sweet Almonds upon it. Others have contrived

3. The Horizontal or Rectangular Barometer (*Fig. 30.*) hermetically sealed at A, and filled with Mercury from D to E; then as the upper Surface of it rises in the Tube, suppose from E to F, the lower will be driven from D to G, as many Times farther, as this Part of the Tube is less than that at E. But it often happens, that some Parts of the Mercury break off from the rest in the Leg BC, and are left behind. This Inconvenience is remedied in

4. The Diagonal Barometer ABC (*Fig. 31.*) wherein the Mercury, instead of rising from B to D (suppose that Space to correspond to the Scale of Variation in a strait Tube) will rise from B to A; for it will always stand at the same perpendicular Height, whatever be the Inclination of the Tube; because Fluids press only according to their perpendicular Altitude *. But the Tube AB must not be too much inclined, lest the Mercury break in it, as in the former.

* Chapter I. §. 9.

5. AB (*Fig. 32.*) is Dr. Rook's Wheel-Barometer, wherein ABD is a Tube filled with Mercury from *a* to E; *a* is an Iron Ball, swimming on the Surface of the Mercury; this as it subsides with the Surface of the Mercury, draws the little Wheel *m n* round, to whose Circumference it is fixed by Means of the String *ac* *: This Wheel carries the Index PQ, which points to the graduated Edge of the Circle KL, and by its Motion shews the most minute Variations of the Mercury. When the Ball *a* is raised by the Mercury on which it swims, the Index is drawn the contrary Way by a lesser Ball *b*, which hangs on the other Side the Wheel. The Friction in this Machine, unless it be made with great Accuracy indeed, renders it useless.

6. The pendent Barometer is another Contrivance to render the Variations more sensible. It consists of a small conical Tube, (represented *Fig. 33.*) hermetically sealed at A, and filled with Mercury from C to D, and empty from thence to A. Now, supposing the Gravity of the Air encreased, it will raise the Mercury higher in the Tube, and so force it into a narrower Part; by which Means the Column becoming longer, its perpendicular Pressure upon the Air below will be proportionably encreased.

* The Tube is smaller at *a* than at E, that the greatest Variation may be at that Surface of the Mercury on which the Ball rests.

Differt. IV. *Of the Barometer.* 103

On the contrary, when the Air becomes lighter, the Mercury descends into a larger Part of the Tube, and by that Means has the Length of its Column proportionably contracted. The Inconvenience that attends this Barometer, is that the Tube must be very small, otherwise the Mercury will fall out; or the Air will be apt to get into it, and divide the Column in several Places; and when the Tube is very small, the Friction of the Mercury against the Sides of it, will hinder it from rising and falling freely.

7. Dr. *Hook*, observing how unfit the common Barometer was to be used on Board of Ship, by Reason its Position ought to be steady, contrived the following one, called, from its Use, a *Marine* Barometer, consisting of two Parts, the one *AB* (*Fig. 34.*) the common Spirit Thermometer, the other *CD*, a Tube filled with Air from *C* to *E*, and from thence to the End *D* with tinged Water. This End is immersed in the same Fluid contained in the Vessel *GF*; and having its Surface exposed to the Pressure of the external Air. Now, the last of these Machines will be affected both by the Warmth of the external Air, and also by its Pressure: The former dilating the Air included in *CE*, and by that Means driving the Water downwards; the latter pressing it up higher in the Tube: Whereas the other, *viz.* *AB*, is affected by the Warmth of the Air alone. Consequently,

sequently, were these Instruments graduated in such a Manner, that, if the Gravity of the external Air should always remain the same it was, when the Instruments were made, their Variations (then only depending on its Warmth) should exactly correspond with each other; that is, when the Spirit in the Tube A B, should ascend to 1, the Water in C D, should descend to 1, &c. Then, whenever their Variations should be observed to differ from each other, the Difference could only be ascribed to some Alteration in the Pressure of the Air upon the Surface of the Water in the Vessel G F. In Proportion therefore as this Difference is greater, or less, so is the Alteration in the Gravity of the Air, from what it was when the Instruments were adjusted. For Instance, when the Water stands above the Division, which corresponds to that, which the Spirit points to in the other Machine, it is an Indication, that the Pressure of the Air is greater at that Time, than when the Instruments were graduated, and *vice versa*.

This Machine is not only more useful at Sea, than the common one, as not requiring a steady Position; but may have its Scale of Variation considerably enlarged, by making the Bore of the Tube C D very small, in Proportion to the Capacity of its Head C.

But

Differt. IV. *Of the Barometer.* 105

But it is observed, that in long keeping this Instrument, the included Air loses somewhat of its Elasticity ; whereby, in Process of Time, the Water stands higher than it ought, and therefore indicates the Gravity of the Air to be greater than what it is.

In the Philosophical Transactions N^o. 427. I have given an Account of a Barometer, wherein the Scale of Variation may be encreased *ad Infinitum*. The Description of it is as follows: A B C D (*Fig. 35.*) is a cylindrical Vessel, filled with a Fluid to the Height W, in which is immersed the Barometer S V, consisting of the following Parts: The Principal of which is a Glass Tube T P (represented separately at *t p*) whose upper End T is hermetically sealed: This End does not appear to the Eye, being received into the lower End of a Tin Pipe G H, which in its other End G receives a cylindrical Rod, or Tube ST, and thereby fixes it to the Tube TP. This Rod ST may be taken off, in Order to put in its stead a larger, or lesser, as Occasion requires. S is a Star at the Top of the Rod ST, and serves as an Index, by pointing to the graduated Scale L A, which is fixed to the Cover of the Vessel A B C D. MN is a large cylindrical Tube made of Tin (represented separately at *m n*) which receives in its Cavity the smaller Part of the Tube TP, and is well cemented to it
at

But

at both Ends, that none of the Fluid may get in.

The Tube TP, with this Apparatus, being filled with Mercury, and plunged into the Bason V, which hangs by two, or more Wires, upon the lower End of the Tube MN, must be so poized, as to float in the Liquor contained in the Vessel A B C D, and then the whole Machine will rise, when the Atmosphere becomes lighter, and *vice versa*.

I shall here add a Computation, in Order to shew the Possibility of the Variation being infinite, upon a given finite Variation of the Weight of the Atmosphere, and withal, the Reason why it may be so. And for the Sake of those who would see a Mathematical Proof of it, I shall give the Demonstration in a Note below. *

Let

* Let the specific Gravity of Quicksilver be to that of Water, or to the Liquor the Barometer floats in, as s to 1 ; and if it be proposed, that the Variations in this compound Barometer shall be to the contemporary Variations of the common Barometer in the given Ratio of n to 1 , this Effect will be obtained, by making the Diameter of the Rod ST to the Diameter of the Cavity of the

Tube HI, as $\sqrt{\frac{n+s}{ns}}$ to 1 , which may be thus demonstrated.

Let us suppose, that the Variation in the Height of the Quick-silver in the common Barometer, which we call v , is such, that a cubic Inch of Quicksilver shall rise into the Vacuum XT; in Order to which, a cubic Inch of Quicksilver must rise from the Vessel V; that is, the Surface P must subside so far, that a cubic Inch of Water (if that be the Fluid made Use of) shall enter the Vessel V, by which Means the Barometer with the Parts annexed will be heavier by a cubic Inch of the Fluid. Now

II.

get

ing

the

ires,

must

con-

the

here

er to

being

the

the

Sake

atical

ation

Let

Water,

if it be

shall be

r in the

king the

of the

rated.

e Quick-

ch, that

XT; in

from the

at a cubic

enter the

s annexed

Now

Fig. 28. *Fig.* 29. *P.* 100. *Fig.* 30. *Fig.* 31. *P.* 101.





Fig. 32. *P.* 102. *Fig.* 33.






Fig. 34. *P.* 103.





Differ

Let
of is V
the W

Now
make the
drostatics)
above the
such a M
nitude of
Square of
at H (the
Bases.) T
rometer w
the comm
their Base

But, by
the stagna
mented b
compound
Column o
silver (to v
by a Colu
second Af
were the h
be lighter

on this A

But now
cury was a

to be $d \cdot v$;

scnt of th
 ddd

scnt —

$s s$
fore the w
cent of the

$d d$
as $d +$ —

s
cause the T

Differt. IV. *Of the Barometer.* 107

Let it be supposed, that the Fluid made use of is Water, and that the given Variation in the Weight of the Atmosphere is such, that,
by

Now this additional Weight of a cubic Inch of Fluid, will make the whole Barometer subside (according to the Laws of Hydrostatics) till a cubic Inch of the Rod HS, immediately extant above the Surface at W, shall come under it; but the Length of such a Magnitude of HS will exceed the Length of an equal Magnitude of Quicksilver in the larger Tube X, as many Times as the Square of the Diameter at X exceeds the Square of the Diameter at H (the Lengths of equal Cylinders being reciprocal to their Bases.) That is, the perpendicular Descent of the compound Barometer will be to v , the perpendicular Ascent of the Mercury in the common Barometer, as d to 1 (supposing this the Ratio of their Bases, and consequently will be equal to dv).

But, by this Descent, the Distance PW, between the Surface of the stagnant Quicksilver and the Top of the Fluid, will be augmented by a Column, whose Height is dv , the Descent of the compound Barometer; and consequently the Weight of the whole Column of the Fluid pressing on the lower Surface of the Quicksilver (to which the Height X is partly owing) will be encreased by a Column of that Length; and this Increase would produce a second Ascent of the Mercury at X equal to itself, namely, dv , were the Fluid as heavy as Quicksilver; but since it is supposed to be lighter in the Ratio of s to 1, the Ascent of the Quicksilver,

on this Account, will only be $\frac{dv}{s}$.

But now, as in the former Case, when the Ascent of the Mercury was v , the Descent of the compound Barometer was shewn

to be dv ; so here, the Ascent of the Mercury being $\frac{dv}{s}$ the Descent of the common Barometer will be $\frac{ddv}{s}$ and the next Descent $\frac{dddv}{ss}$ and the next $\frac{d^4v}{s^3}$ and so on to Infinity. There-

fore the whole Descent of the compound Barometer, is to the Ascent of the Mercury in the common Barometer, that is, n is to 1

as $d + \frac{dd}{s} + \frac{ddd}{ss} + \frac{d^4}{s^3} + \dots$ to 1, or as $\frac{ds}{s-d}$ to 1; be-

cause the Terms of the Series being in geometrical Proportion, the
Sum

by pressing upon the Surface of it at W, the Surface of the Mercury at X may be raised an Inch higher (measuring from its Surface at P) than before; and that the Breadth of the Cavity of the Tube at X, and of the Basin at P are such, that by this Ascent of the Mercury, there may be a cubic Inch of it in the Cavity X more than before, and consequently in the Basin a cubic Inch less. Now, upon this Supposition, there will be a cubic Inch of Water in the Basin more than there was before; because the Water will succeed the Mercury to fill up its Place. Upon this Account, the whole Machine will be render'd heavier, than it was before, by the Weight of a cubic Inch of Water, and therefore will sink, according to the Laws of Hydrostatics (Chap II. §. 5.) till a cubic Inch of that Part of the Rod WS, which

Sum of them all is $\frac{ds}{s-d}$ Hence we have $n = \frac{ds}{s-d}$ and therefore $ns = ds + dn$; that is, $1 : d :: n + s : ns :: \frac{n+s}{ns} : 1$ and therefore, by extracting the Square Roots of each Term in the Proportion, $1 : \sqrt{d}$ (that is, the Diameter of ST to the Diameter of HI) as $\sqrt{\frac{n+s}{ns}}$ to 1. Q. E. D.

Example 1. Putting $s=14$ and $n=1$, the Variation in each Barometer will be equal, by taking the Diameter of ST to the Diameter of HI, as $\sqrt{\frac{15}{14}}$ to 1, that is, as 30 to 29 nearly.

Example 2. If n be put infinite, the Diameter of ST will be to the Diameter of HI, as $\sqrt{\frac{1}{s}}$ to 1, or 1 to $\sqrt{14}$; that is, as 1 to $3\frac{1}{4}$ nearly.

was above the Surface of the Water at W, comes under it. Then, if we suppose this Rod so small, that a cubic Inch of it shall be fourteen Inches in Length, the whole Machine will sink fourteen Inches lower into the Fluid than before, and consequently the Surface of the Mercury in the Bason will be pressed more than it was before, by a Column of Water fourteen Inches high. But the Pressure of fourteen Inches of Water is equivalent to one of Mercury (because Water is about fourteen Times lighter than Mercury) this additional Pressure therefore will make the Mercury ascend at X, as much as the supposed Variation in the Weight of the Air did at first. This Ascent will give Room for a second cubic Inch of Water to enter the Bason; the Machine will therefore be again rendered heavier by the Weight of so much Water, and accordingly will subside fourteen Inches farther. This will occasion another additional Pressure of Water, which will raise another Inch of Mercury, and make the Machine sink fourteen Inches more, and so on, without ever approaching nearer to an *Æquilibrium* with the external Air: and therefore a Scale, answering to the Variation of this Barometer, ought strictly and properly to be of an infinite Length; because after this Barometer has sunk or risen thousands of Miles (if that were possible) it would still have the same Tendency to sink or rise on, as when it first set out.

Now, was the Rod WS so small, that a cubic Inch of it should be more than fourteen Inches long (the other Parts remaining as was supposed above) the Variation in this Barometer would be more than infinite, or negative with Respect to those of the common Barometer. The Meaning of which is, that whereas in the common Barometer, the suspended Column of Mercury, by its rising or falling, approaches nearer to an *Æquilibrium* with the external Air, this Barometer would continually recede from an *Æquilibrium* with it; so that the farther it should move up or down, instead of acquiring by that Means a less Tendency to move on, as the Mercury in the common Barometer does, it would acquire a greater.

On the contrary, when a cubic Inch of the Rod is less than fourteen Inches in Length, the Variation will be finite; and may be made to bear any Proportion to those of the common Barometer whatever, as demonstrated in the foregoing Note.

While I am writing this, another Method occurs to me of making a Barometer, wherein the Scale of Variation shall bear any Proportion to that of the common one. It is this; Let there be a compound Tube, as ABC (*Fig. 36.*) hermetically sealed at A, and open at C, empty from A to D, filled with Mercury from thence to B, and from thence to E with Water;

Differt. IV. *Of the Barometer.* III

ter: Then, if the Tube FC be a little more than five Times less in Diameter than the Tube FA, the Variation in the lower Surface of the Water at E will be infinite; if it be above so many Times less, it will be more than infinite, otherwise it will be finite. See the Demonstration in the Note *.

That

* Let v denote a given Variation in the common Barometer, x the correspondent Variation at E sought. Let the Ratio of m to 1, express that of the specific Gravity of Mercury, to that of Water; and d to 1, that of the Diameter of the Tube FA to FC. Then the Variation at E, the lower Surface of the Water, being supposed x , the Variation of it at B, the upper Surface of it will

be $\frac{x}{dd}$ and consequently GE, the Difference of the Legs EK and KB, will vary $x + \frac{x}{dd}$. Again, the Variation of the Surface of

the Mercury at B will be the same with that of the Water in the same Place, viz. $\frac{x}{dd}$; and, if the Tube is of the same Diameter

at D, as at B, the Variation of the Surface at D will also be the same, that is, $\frac{x}{dd}$: The Sum of both Variations, or the Variation

of HD the Difference of the Legs, will therefore be $\frac{2x}{dd}$. Now the Pressure of the Mercury and Water together upon the Air at E, is owing to the Lengths of HD and GE; and since one of these will always shorten, when the other lengthens, the Variation in their Pressure will depend on the Variation of the Difference of their Weights, that is, of the Difference between the Weight of $x + \frac{x}{dd}$ and of $\frac{2x}{dd}$. But the Weight of $x + \frac{x}{dd}$ (being the Weight of a Column of Water) compared to that of a Column of Mercury of the same Length, is only $\frac{x + \frac{x}{dd}}{m}$. The Difference therefore between

That the Variation in this Barometer may be infinite, will appear from the following Computation.

Let the Proportion between the Bores of the Tube AF and FC be such, that when HD, the Difference of the Legs wherein the Mercury is contained, is augmented one Inch, GE the Difference of the Legs, wherein the Water is contained, shall be diminished fourteen; then, as much as the Pressure of the Mercury is augmented, that of the Water will be diminished, and so the Pressure of both taken together will remain as it was. And consequently, after it has began to rise, it will always have the same Tendency to rise on, without ever coming to an *Æquilibrium* with the Air.

How far this Barometer will succeed in Practice, must be left to Experience to determine.

between $\frac{x + \frac{x}{dd}}{m}$ and $\frac{2x}{dd}$ will always be equal to the Variation in

the common Barometer, and therefore $\frac{2x}{dd} \frac{x + \frac{x}{dd}}{m} = v$, and by

the common Method of Reduction, $x = \frac{v m d d}{2m - dd - 1}$: That is, x :

$v :: m d d : 2m - dd - 1$. Now, if we put $m=14$, and $d=5$, $2m - dd - 1$ will be as much as $2m$, and therefore $2m - dd - 1$ will be equal to nothing; and so x being by the Proportion as many Times more than v , as $m d d$ is than nothing, 'tis infinite. And if m be put $=14$, and $d=5$, $m d d$ will be equal to 350, and $2m - dd - 1 = 2$; and therefore the Variations, in this Case, will be to those in the common Barometer, as 175 to one.

Pro-

Differt. IV. *Of the Barometer.* 113

Probably, if the Bore of the Tube FC be made very small, *viz.* about the twentieth Part of an Inch Diameter, the Air will not ascend through the Water, as it is apt to do through the Mercury in the pendent Barometer; and the Smallness of the Bore will not prevent the Water, from moving, near so much as it does the Mercury in that Barometer.

There is an Improvement of another Kind in the common Barometer, whereby it is rendered *portable*. The Tube containing the Mercury, instead of having its lower End immersed in a Vessel of that Fluid, has it tied up in a leathern Bag, not quite full of Mercury. And though the external Air cannot get into the Bag to suspend the Mercury in the Tube, by pressing on its Surface, as in the common one; yet it has the same Effect by pressing on the Outside of the Bag, which, being pliant, yields to the Pressure, and keeps the Mercury suspended in the Tube at its proper Height. This Bag is generally inclosed in a little Box, through the Bottom of which passes a Screw; with this Screw the Bag may be compressed, so as to force the Mercury up to the Top of the Tube; which keeps it steady, and hinders it from breaking the Tube by dashing against the Top when it is carried about, as it otherwise would be apt to do.

See

See more on the Subject of this Dissertation, *Weidleri Institutiones Mathematicæ*. p. 608. *Melchior Verdries Phys. Pars specialis*, Cap. IV. §. 15. Mr. *Paschal's* Traité de l'Equilibre des Liqueurs. *Sinclair's* Ars magna Gravitatis & Levitatis. *Mariotte's* Second Essay de la Nature de l'Air. *Philosoph. Burgund. Tom. II.* p. 850. *Saul's* Treatise on the Barometer. *Regnault's* Philosoph. Conversat. 22. *Clare's* Motion of Fluids, p. 141. *Mem. de l'Acad.* 1705, 1711. *Philosophical Transactions* N^o. 9, 10, 11, 55, 86, 91, 165, 181, 185, 208, 229, 236, 237, 240, 243, 269, 351, 366, 385, 388, 405, 406, 427. *Cotes's* Hydrostatical and Pneumatical Lecture. Lect. 7. With several other Authors referred to in Mr. *Johnson's* Quæstiones Philosophicæ, Cap. VI. Quæst. 36, 37.



DISSERTATION V.

Of the Origin of the Winds.

THE Wind is no other, than the Motion of the Air, upon the Surface of the Globe. Some of the Ancients took it to be Air, rushing out of the Bowels and Cavities of the Earth: And others thought it an Exhalation from its Surface. But these are *Hypotheses* too chimerical to stand in Need of a particular Confutation. Some of the Moderns, who held a *Plenum*, have accounted for it thus. They imagined, that the Air being confined above, as it must be, if we suppose a *Plenum*, would, when more than ordinarily rarefied, or stocked with Vapours, drive away the neighbouring Air, in order to make room for itself; and by this Means occasion a Wind. Others, observing a constant and perpetual easterly Wind to blow at the *Equator*, ascribed its Origin to the diurnal Rotation of the Earth, about its Axis from West to East; which they thought would occasion the Air upon its Surface, to seem to move the contrary Way, being in some Measure left behind. But, whereas there are Winds, in some Places near the *Equator*,
Q
that

116 *Of the Origin of Winds.* Part II.

that blow on other Points of the Compass (as we shall see hereafter) this *Hypothesis* is insufficient. Besides, the Air pressing upon the Surface of the Earth by its Gravity, like other Bodies; and having nothing to hinder it from moving freely along with it, must necessarily, in Time, acquire an equal Degree of Velocity, and so keep Pace with it, all the Way round.

The principal Cause of the Wind, or, in other Words, of the Air's moving from Place to Place, upon the Surface of the Earth, is the Atmosphere's being heated over one Part more than over another. For, in this Case, the warmer Air, being rendered specifically lighter than the rest, rises up into the superior Parts of the Atmosphere, and there diffuses itself every Way; while the neighbouring inferior Air rushes in from all Parts at the Bottom, to restore the *Æquilibrium*.

Upon this Principle it is, that most of the Winds may be accounted for.

To begin with those which blow under the *Equator*.

1. Under the *Equator*, the Wind is always observed to blow from the East Point*.

For,

* For the Reader's Ease (who perhaps is not furnished with the Philosophical Transactions) I shall here insert by Way of Note, from Dr. *Halley's* Account, so much of the History of the Winds, as may be necessary for the understanding this Theory.

The

Diff. V. Of the Origin of Winds. 117

For, supposing the Sun to continue vertical over some one Place, the Air will be most rarified there; and consequently, the neighbouring

“ The universal Ocean, says he, may most properly be divided into three Parts, viz. 1. The *Atlantic* and *Æthiopic* Seas.
“ 2. The *Indian* Ocean. 3. The great *South* Sea, or the *Pacific* Ocean.

“ I. In the *Atlantic* and *Æthiopic* Seas, between the *Tropics*, there is a general easterly Wind all the Year long, without any considerable Variation; excepting, that it is subject to be deflected therefrom, some few Points of the Compass, towards the North, or South, according to the Position of the Place.

“ 1. Near the Coast of *Africa*, as soon as you have passed the *Canary* Isles, you are sure to meet a fresh Gale of North-east Wind, about the Latitude of twenty-eight Degrees North; which seldom comes to the Eastwards of the East-north-east, or passes the North-north-east. This Wind accompanies those bound to the Southward, to the Latitude of ten Degrees North, and about an hundred Leagues from the *Guinea* Coast; where, till the fourth Degree of North Latitude, they fall into *Calms* and *Tornadoes*, or sudden Storms.

“ 2. Those bound to the *Caribbe* Isles, find, as they approach the *American* Side, that the aforesaid North-east Wind becomes still more and more easterly, so as sometimes to be East, sometimes East by South, but yet most commonly to the Northward of the East, a Point or two, seldom more. 'Tis likewise observed, that the Strength of these does gradually decrease, as you sail to the Westward.

“ 3. That the Limits of the *Trade* and *variable* Winds in this Ocean, are farther extended on the *American* Side, than the *African*; for, whereas you meet not with this certain Wind, till after you have pass'd the Latitude of twenty eight Degrees on this Side; on the contrary Side it commonly holds to thirty, thirty-one, or thirty-two Degrees of Latitude; and this is verified likewise to the Southward of the *Æquinoctial*; for near the *Cape of Good Hope*, the Limits of the *Trade* Winds are three or four Degrees nearer the *Line*, than on the Coast of *Brazil*.

Q 2

“ 4. That

118 *Of the Origin of Winds.* Part II.

ing Air will rush in from every Quarter with equal Force. But, as the Sun is continually shifting to the Westward, the Part, where the
Air

“ 4. That from the Latitude of four Degrees North, to the aforefaid Limits on the South Side of the *Equator*, the Winds are generally and perpetually between the South and East, and most commonly between the South-east and East; observing always this Rule, that on the *African* Side, they are more southerly, on the *Brazilian* more easterly, so as to become almost due East, the little Deflection they have being still to the Southward. In this Part of the Ocean, it has been my Fortune to pass a full Year, in an Employment that obliged me to regard more than ordinarily the Weather, and I found the Winds constantly about the South-east, the most usual Point South-east by East: When it was easterly, it generally blew hard, and was gloomy, dark, and sometimes rainy Weather: If it came to the Southwards, it was generally serene, and a small Gale next to a Calm; but this not very common. But I never saw it to the Westwards of the South, or Northwards of the East.

“ 5. That the Season of the Year has some small Effect on these *Trade* Winds; for that when the Sun is considerably to the Northward of the *Equator*, the South-east Winds, especially in the Streight of this Ocean (if I may so call it) between *Brazil*, and the Coast of *Guinea*, do vary a Point or two to the Southward, and the North-east become more easterly; and, on the contrary, when the Sun is towards the *Tropic of Capricorn*, the South-easterly Winds become more easterly, and the North-easterly Winds, on this Side the *Line*, veer more to the Northward.

“ 6. That as there is no general Rule, that admits not of some Exception, so there is in this Ocean a Tract of Sea, wherein the southerly and South west Winds are perpetual, viz. all along the Coast of *Guinea*, for above five hundred Leagues together, from *Sierra Leona*, to the Isle of *St. Thomas*: For the South-east *Trade* Wind having pass'd the *Line*, and approaching the Coast of *Guinea* within eighty or an hundred Leagues, inclines towards the Shore, and becomes South-south-east; and by Degrees, as you come nearer, it veers about to South, South-south-west, and in with the Land South-west, and
“ sometimes

Diff. V. Of the Origin of Winds. 119

Air is most rarefied, is carried the same Way :
and therefore the Tendency of all the lower Air,
taken together, is greater that Way, than any
other.

“ sometimes West-south-west. These are the Winds, which are
“ observed on this Coast when it blows true ; but there are fre-
“ quent Calms, violent sudden Gusts, called *Tornadoes*, from all
“ Points of the Compass, and sometimes unwholsome foggy
“ easterly Winds, called *Hermitæ*, by the Natives, which too
“ often infest the Navigation of these Parts.

“ 7. That to the Northwards of the *Line*, between four and
“ ten Degrees of Latitude, and between the Meridians of *Cape*
“ *Verde*, and of the eastermost Islands that bear that Name, there
“ is a Tract of Sea, wherein it were improper to say, there is
“ any *Trade Wind*, or yet a *variable* ; for it seems condemned to
“ perpetual Calms, attended with terrible Thunder and Light-
“ ning, and Rains so frequent, that our Navigators from thence
“ call this Part of the Sea, the *Rains* : The little Winds that
“ are, being only some certain Gusts, of very little Conti-
“ nuance, and less Extent ; so that sometimes each Hour you
“ shall have a different Gale, which dies away into a Calm
“ before another succeeds : And in a Fleet of Ships in Sight of
“ one another, each shall have the Wind from a several Point of
“ the Compass : With these weak Breezes, Ships are obliged to
“ make the best of their Way to the Southward, through the
“ aforesaid six Degrees ; wherein it is reported some have been
“ detained whole Months for want of Wind. —

“ II. In the *Indian Ocean*, the Winds are partly general,
“ as in the *Æthiopic Ocean* ; partly periodical, that is, half
“ the Year they blow one Way, and the other half, near upon
“ the opposite Points ; and these Points and Times of shifting,
“ are different in different Parts of this Ocean.

“ 1. Between the Latitudes of ten Degrees and thirty Degrees
“ South, between *Madagascar* and *Hollandia nova*, the *Gene-*
“ *ral Trade-Winds* about the South-east and by East, are found
“ to blow all the Year long, to all Intents and Purposes,
“ after the same Manner, as in the same Latitudes in the
“ *Æthiopic Ocean*, as it is described in the fourth Remark
“ aforesaid.

“ 2. That

120 *Of the Origin of Winds.* Part II.

other. Thus the Tendency of the Air towards the West, becomes general, and its Parts impelling one another, and continuing to move

“ 2. That the aforesaid South-east Winds extend to within
 “ two Degrees of the *Equator*, during the Months of *June*,
 “ *July*, and *August*, &c. to *November*, at which Time, between
 “ the South Latitude of three and ten Degrees, being near the
 “ *Meridian* of the North End of *Madagascar*, and between two
 “ and twelve South Latitude, being near *Sumatra* and *Java*;
 “ the contrary Winds from the North-west, or between the
 “ North and West, set in, and blow for half a Year, viz.
 “ from the Beginning of *December* till *May*: And this *Mon-*
 “ *soon* is observed as far as the *Molucca* Isles.

“ 3. That to the Northward of three Degrees South Lat-
 “ tude, over the whole *Arabian* and *Indian* Sea, and Gulf of
 “ *Bengal*, from *Sumatra* to the Coast of *Africa*, there is another
 “ *Monsoon*, blowing from *October* to *April*, upon the North-
 “ east Points: But in the other half Year, from *April* to *October*,
 “ upon the opposite Points of South-west and West-south-west,
 “ and that with rather more Force than the other, accompanied
 “ with dark, rainy Weather; whereas the North-east blows clear.
 “ 'Tis likewise to be noted, that the Winds are not so constant,
 “ either in Strength or Point, in the Gulf of *Bengal*, as they
 “ are in the *Indian* Sea, where a certain steady Gale scarce ever
 “ fails. 'Tis also remarkable, that the South-west Winds, in
 “ these Seas, are generally more southerly on the *African* Side,
 “ and more westerly on the *Indian*.

“ 4. There is a Tract of Sea to the Southwards of the
 “ *Equator*, subject to the same Changes of the Winds, viz.
 “ near the *African* Coast, between it and the Island *Madagascar*,
 “ or *St. Laurence*, and from thence Northwards, as far as the
 “ *Line*; wherein from *April* to *October*, there is found a con-
 “ stant fresh South-south-west Wind, which, as you go more
 “ northerly, becomes still more and more westerly, so as to fall
 “ in with the West-south-west Winds, mentioned before in those
 “ Months of the Year to be certain to the Northward of the
 “ *Equator*. What Winds blow in those Seas, for the other
 “ half Year, I have not yet been able to obtain to my full Satis-
 “ faction: The Account which has been given me, is only this,
 “ That

Diff. V. *Of the Origin of Winds.* 121

move till the next Return of the Sun, so much of its Motion as was lost by his Absence, is again restored, and therefore the easterly Wind becomes *Perpetual*.

Some

“ That the Winds are much easterly hereabouts, and as often to
“ the North of the true East, as to the Southward thereof.

“ 5. That to the Eastward of *Sumatra* and *Malacca*, to the
“ Northwards of the *Line*; and along the Coast of *Camboia*
“ and *China*, the *Monsoons* blow North and South; that is to
“ say, the North-east Winds are much northerly, and the
“ South-west much southerly. This Constitution reaches to the
“ Eastward of the *Philippine* Isles, and as far northerly as
“ *Japan*; the northern *Monsoon* setting in, in these Seas, in
“ *October* or *November*; and the southern in *May*, blowing all
“ the Summer Months. Here it is to be noted, that the Points of
“ the Compass from whence the Winds come, in these Parts of
“ the World, are not so fixed, as those lately described; for the
“ southerly will frequently pass a Point or two to the Eastwards
“ of the South, and the northerly as much as to the Westwards
“ of the North, which seems occasioned by the great Quantity
“ of Land which is interpersed in these Seas.

“ 6. That in the same Meridians, but to the Southwards of
“ the *Equator*, being that Tract lying between *Sumatra* and
“ *Java* to the West, and *New Guinea* to the East, the same
“ northerly and southerly *Monsoons* are observed; but with this
“ Difference, that the Inclination of the northerly is towards
“ the North-west, and of the southerly towards the South-east:
“ But the *Plagæ Venti* are not more constant here than in the
“ former, *viz. variable* five or six Points. Besides, the Times of
“ the Change of these Winds are not the same, as in the *Chinese*
“ Seas, but about a Month, or six Weeks later.

“ 7. That the contrary Winds do not shift all at once, but in
“ some Places the Time of the Change is attended with *Calms*,
“ in others with *variable* Winds; and it is particularly remark-
“ able, that the End of the westerly *Monsoon*, in the Seas of
“ *China*, are very subject to be tempestuous. The Violence of
“ these Storms is such, that they seem to be of the Nature of the
“ *West-India* Hurricanes, and render the Navigation of these Parts
“ very unsafe about that Time of the Year. These Tempests are
“ by

122 *Of the Origin of Winds.* Part II.

Some are inclined to think, that the continual shifting of the Sun to the Westward, should produce a westerly Wind under the *Equator*, by causing the Current of the Air from the West to exceed and over-balance that,

“ by our Seamen, usually term’d the *Breaking up of the Mon-*
“ *soons.*

“ III. The Third Ocean, called *Mare Pacificum*, whose Ex-
“ tent is equal to that of the other two (it being from the West
“ Coast of *America* to the *Philippine* Islands, not less than an hun-
“ dred and fifty Degrees of Longitude) is that which is least
“ known to our own, or the neighbouring Nations : That Navi-
“ gation that there is on it, is by the *Spaniards*; who go
“ yearly from the Coast of *New-Spain* to the *Manilha*’s: But
“ that but by one beaten Track; so that I cannot be so particular
“ here, as in the other Two. What the *Spanish* Authors say of
“ the Winds, they find in their Courses; and what is confirmed
“ by the old Accounts of *Drake* and *Candish*, and since by
“ *Schooten*, who sailed the whole Breadth of this Sea, in the
“ southern Latitude of fifteen or sixteen Degrees, is, that there is
“ a great Conformity between the Winds of this Sea, and those
“ of the *Atlantic* and *Ethiopic*; that is to say, that to the North-
“ ward of the *Equator*, the predominant Wind is between
“ the East and North-east; and to the Southwards thereof, there
“ is a constant, steady Gale, between the East and South-east,
“ and that on both Sides the *Line*, with so much Constancy, that
“ they scarce ever need to attend the Sails; and strength, that it
“ is rare to fail of crossing this vast Ocean in ten Weeks
“ Time; which is about an hundred and thirty Miles a Day;
“ Besides, ’tis said, that Storms and Tempests are never known
“ in these Parts, wherefore some have thought it might be as
“ short a Voyage to *Japan* and *China*, to go by the Streights of
“ *Magellan*, as by the Cape of *Good Hope*.

“ The Limits of these General Winds are much the same
“ as in the *Atlantic* Sea, viz. about the thirtieth Degree of La-
“ titude on both Sides. Besides, a farther Analogy between the
“ Winds of this Ocean, and the *Ethiopic*, appears in that,
“ that upon the Coasts of *Peru*, they are always much southerly,
“ like as they are found near the Shores of *Angola*.

which

Diff. V. Of *the Origin of Winds.* 123

which opposes it from the East. For, whereas the eastern Air retains its Heat some time after the Sun is removed from it; it must always be rarefied to a greater Degree, and also to a greater Distance from the Place to which the Sun is vertical, than the western Air is; and therefore the western Air, being more ponderous, should be an Over-balance for the eastern, and drive its Current before it.

But it is to be observed, that we are not to consider the Point to which the Sun is vertical, but the Point of greatest Rarefaction (which, upon Account of the Sun's Motion to the Westward, lies always to the Eastward); and then see, which Side of the Column of Air, incumbent over that Point, sustains the greater Pressure from the neighbouring Columns. Now, although the Air is rarefied even farther to the East of this Point, than to the West, yet still, if we suppose the Point to keep its place, the Air over it will sustain an equal Degree of Pressure on each Side. For, since no Column can be assigned on the western Side, but one also on the eastern, may be found under an equal Degree of Rarefaction, and therefore of the same specific Gravity: And since Fluids of equal Heights, and the same specific Gravities (whatever be the Breadth of their Columns) press equally against equal Objects, (Chap. I. §. 9.) 'tis very evident, that the Column of Air, over the

R

Point

124 *Of the Origin of Winds.* Part II.

Point of greatest Rarefaction, is pressed equally both Ways; and so, upon this Supposition, each Wind will blow towards that Point with equal Force. But, if we suppose the Point of greatest Rarefaction to shift towards the West, we shall find, that this *Æquilibrium* will by that Means be destroyed, and the Motion of the Air (upon the whole) determined that Way. For let us consider any Portion of the western Air approaching towards the Point of greatest Rarefaction, if that Point shifts, in the mean Time, towards the West, then will that Portion reach the said Point sooner than it otherwise would have done; thereby losing a Part of its Motion, by which Means the westerly Current will be diminish'd. Again, if, while the East Wind blows towards the Point of greatest Rarefaction, that Point moves on before it, then will the eastern Air have a greater Quantity of Motion, than it otherwise would have had; that, which should have been an Impediment to it, being, upon this Supposition, in some Measure withdrawn; and so the East Wind will be augmented. Thus, the West Wind having its Force lessen'd by the Motion of the Sun, and the East one being increased, the latter at length as it were absorbs the former, and carries it away in its own Direction.

2. On each Side of the *Equator*, to about the thirtieth Degree of Latitude, the Wind is found

Diff. V. *Of the Origin of Winds.* 125

found to vary from the East Point, so as to become North-east on the northern Side, and South-east on the southern.

The Reason of which is, that, as the *equatorial* Parts are hotter than any other, both the northern and southern Air, ought to have a Tendency that Way; the northern Current therefore, meeting in this Passage with the eastern, produces a North-east Wind on that Side; as the southern Current joining with the same, on the other Side the *Equator*, forms a South-east Wind there.

These two Propositions are to be understood of open Seas, and of such Parts of them as are distant from the Land; for near the Shores, where the neighbouring Air is much rarefied, by the Reflection of the Sun's Heat from the Land, it frequently happens otherwise particularly.

3. On the *Guinea* Coast, the Wind always sets in upon the Land, blowing westerly instead of easterly. This is because the Deserts of *Africa* lying near the *Equator*, and being a very sandy Soil, reflect a great Degree of Heat into the Air above them. This therefore being rendered lighter, than that which is over the Sea, the Wind continually rushes in upon the Land to restore the *Æquilibrium*.

4. That Part of the Ocean, which is called the *Rains*, is attended with perpetual Calms, the Wind scarce blowing sensibly either one

126 *Of the Origin of Winds.* Part. II.

Way or other. (See its Situation described in Note, Page 119, Remark 7th). For this Tract being placed between the westerly Wind blowing from thence towards the Coast of *Guinea*; and the easterly Wind blowing from the same Place to the Westward thereof, the Air stands in *Æquilibrio* between both, and its Gravity is so much diminished thereby, that it is not able to support the Vapour it contains, but lets it fall in continual Rain, from whence this Part of the Ocean has its Name.

5. There is a Species of Winds, observable in some Places within the *Tropics*, called by the Sailors *Monsoons*, which, during six Months of the Year, blow one Way; and the remaining six, the contrary.

The Occasion of them in general is this: When the Sun approaches the northern *Tropic*, there are several Countries, as *Arabia*, *Persia*, *India*, &c. which become hotter, and reflect more Heat than the Seas beyond the *Equator*, which the Sun has left; the Winds therefore, instead of blowing from thence to the Parts under the *Equator*, blow the contrary Way; and when the Sun leaves those Countries, and draws near the other *Tropic*, the Winds turn about, and blow on the opposite Point of the Compass.

The Regularity of these Winds making them more than ordinarily useful in Navigation, they are from thence called *Trade Winds*.

Diff. V. *Of the Origin of Winds.* 127

Winds. As to other Circumstances relating to them, and the particular Times and Places of the *Monsoons*, &c. See the Historical Account in the foregoing Note; all which might easily be solved upon the same Principle, had we *Data* to go upon, and were all the several Circumstances of Situation, Heat, Cold, &c. sufficiently known *.

From the Solution of the general *Trade Winds*, we may see the Reason, why in the *Atlantic Ocean*, a little on this Side the thirtieth Degree of North Latitude, or thereabouts, as was observed in the foregoing Dissertation, there is generally a West, or South-west Wind. For, as the inferior Air, within the Limits of those Winds, is constantly rushing towards the *Equator*, from the North-east Point, or thereabouts, the superior Air moves the contrary Way; and therefore after it has reached these Limits, and meets with Air, that has little or no Tendency to

* Some have thought, that the Regularity of the general *Trade Winds* is partly owing to the diurnal Motion of the Moon from East to West. For, as the Sun renders the Air specifically lighter by its Heat, so does the Moon by attracting it, in the same Manner, as it does the Sea, in raising the Tides. But it is to be observed, that as the Moon acts with the greatest Force upon the superior Parts of the Air, and as those superior Parts are unconfined, and therefore more at Liberty to rush in, in Order to restore the *Æquilibrium*; it will from hence follow, that the rushing in of the superior Parts of the Atmosphere will principally contribute towards restoring the *Æquilibrium*; and so the Motion, produced below, will be very inconsiderable.

any

128 *Of the Origin of Winds.* Part. II.

any one Point more than to another, by Reason of the Sun's Distance, it will determine it to move in the same Direction with itself.

In our own Climate we frequently experience, in calm Weather, gentle Breezes blowing from the Sea to the Land, in the Heat of the Day; which *Phænomenon* is very agreeable to the Principle laid down above: For the inferior Air over the Land being rarefied by the Beams of the Sun, reflected from its Surface, more than that which impends over the Water which reflects fewer, the latter is constantly moving on to the Shore, in order to restore the *Æquilibrium*, when not disturbed by stronger Winds from another Quarter *.

From what has been observed, nothing is more easy than to see, why the northern and southern Parts of the World, beyond the Limits of the *Trade Winds*, are subject to such

- * In Confirmation of this, we have an easy, and very pertinent Experiment, related by Mr. *Clare*, in his *Motion of Fluids*.
 “ Take, *says he*, a large Dish, fill it with cold Water; into the
 “ Middle of this put a Water-Plate, filled with warm Water.
 “ The first will represent the Ocean; and the other an Island,
 “ rarefying the Air above it. Blow out a Wax Candle, and if
 “ the Place be still, on applying it successively to every Side of
 “ the Dish, the fuliginous Particles of the Smoak, being visiole
 “ and very light, will be seen to move towards the Plate,
 “ and rising over it, point out the Course of the Air from Sea
 “ to Land. Again, if the ambient Water be warmed, and the
 “ Plate filled with cold Water, let the smoaking Wick of the
 “ Candle be held over the Plate, and the contrary will happen.

Variety

Diff. V. *Of the Origin of Winds.* 129

Variety of Winds. For the Air, upon Account of its Distance from the *Equator*, being undetermined to move towards any fixed Point, is continually shifting from Place to Place, in Order to restore the *Æquilibrium*, wherever it is destroyed ; whether by the Heat of the Sun, the rising of Vapours, or Exhalations, the melting of Snow upon the Mountains, and a great Variety of other Circumstances, more than can be easily enumerated.

We are told by Historians, of certain Caves that emit Wind ; if so, it is when the included Air is rarefied by Heat, and therefore rushes out for want of Room ; or, when the Pressure of the external Air, incumbent upon the Mouth of the Cave, is diminished, and so permits the internal Air to dilate itself, and issue out.

For more on this Subject, see *Nieuwentyt's Religious Philosopher.* *Regnault's Philosophical Conversations.* *Clare's Motion of Fluids.* *Martin's Philosophical Grammar.* And the Authors referred to in Mr. *Johnson's Quæstiones Philosoph.* Cap. IV. Quæst. 1. 2.



DISSERTATION VI.

Of the Formation, and Ascent of Vapours, and their Resolution into Rain, Snow, and Hail.

THAT Vapours are raised from off the Surface of Water by the Action of the Sun's Heat, is agreed on by all : But the Manner in which this is done, has ever been a Controversy among Philosophers ; neither is it at this Time sufficiently explain'd by any one.

If we consult a *Cartesian* upon this Matter, he immediately tells us, that, by the Action of the Sun upon the Water, small Particles of the Water, are formed into hollow Spheres, filled with *Materia Subtilis*, and by that Means becoming lighter than an equal Bulk of Air, are easily buoyed up in it. But, as this *Materia Subtilis* is only a Fiction, the Solution is not to be regarded.

Dr. *Nieuwentyt*, and several other Philosophers, who maintain, that Fire is a particular Substance, distinct from other Matter, account for the Formation and Ascent of Vapours thus : They say, that the Rays of the Sun, or Particles of Fire separated from them, adhering to Particles of the Water, make
up

* See
† See
Philosophi
§ By I

Diff. VI. *Ascent of Vapours, &c.* 131

up little Bodies, lighter than an equal Bulk of Air; which therefore, by the Laws of Hydrostatics, will ascend in it, till they come to an Height where the Air is of the same specific Gravity with themselves. And, that Rain is produced by the Separation of the Particles of the Fire from those of the Water; which last, being then left without Support, can no longer be sustained by the Air, but falls down in Drops of Rain *.

This Opinion is liable to the following Difficulties; *First*, Fire has never been yet proved to be a distinct Element, or a particular Substance †; and the Change of Weight in Bodies in chymical Preparations, heretofore thought to arise from the Adhesion of Particles of Fire, is found to proceed from the Adhesion of Particles of Air §.

Secondly, Should the above-mentioned Supposition be allowed, the fiery Particles, which are joined to the watery ones to buoy them up, must be considerably large, or else a very great Number must fix upon a single Particle of Water; and then a Person, being on the Top of an Hill in a Cloud, would be sensible of the Heat, and find the Rain produced from that Vapour, much colder than the Vapour it-

* See *Nieuwentyt's Religious Philosopher*, Contempl. 19.

† See the Authors referred to in *Mr. Johnson's Questions Philosoph.* Cap. I. Quæst. 30.

§ By *Dr. Hales*, in his *vegetable Statics*.

self: whereas the contrary is evident to our Senses; the Tops of Hills, though in the Clouds, being much colder than the Rain which falls below.

Besides, the Manner in which the Particles of Water should be separated from those of the Fire, so as to fall in Rain, is not easily to be conceived.

The most generally received Opinion is, That by the Action of the Sun, on the Surface of the Water, the aqueous Particles become formed into Bubbles, filled with a *Flatus*, or *warm Air*, which renders them specifically lighter than common Air, and makes them rise therein, till they meet with such as is of the same specific Gravity with themselves *. But I ask,

First, How comes the Air in the Bubbles to be specifically lighter than that without, since the Sun's Rays, which act upon the Water from whence they are raised, are equally dense over all its Surface?

Secondly, If it could be possible for rarer Air to be separated from the denser ambient Air, to form the Bubbles (as Bubbles of soaped Water are blown up by warm Air from the Lungs, whilst the ambient Air is colder and denser) what would hinder the external Air from reducing that, which is inclosed in the Bubbles, immediately to the same Degree of

* Philosophical Transactions, N^o. 192.

Diff. VI. *Ascent of Vapours, &c.* 133

Coldness, and specific Gravity with itself; (Cold being readily communicated through such thin Shells of Water). By which means, the Bubbles would become specifically heavier than the circumambient Air, and would no longer be supported therein; but fall down, almost as soon as they were formed?

Thirdly, If we should grant all the rest of the Supposition, yet the following Difficulty will still remain. If Clouds are made up of Bubbles of Water filled with Air, why do not those Bubbles always expand, when the ambient Air is rarefied, and presses less upon them than it did before; and why are they not condensed, when the ambient Air is condensed by the Accumulation of the superior Air? But if this Condensation and Rarefaction should happen to them, the Clouds would always continue at the same Height, contrary to Observation; and we should never have any Rain.

The two last Opinions are more largely examined by Dr. *Desaguliers* in the Philosophical Transactions N^o. 407. After which he endeavours to establish one of his own.

He observes, with Sir *Isaac Newton*, that, when by Heat or Fermentation the Particles of a Body are separated from their Contact, their repulsive Force grows stronger, and the Particles exert that Force at greater Distances; so that the same Body shall be expanded into a very large Space, by becoming fluid; and

may sometimes take up more than a Million of Times the Room it did in a solid and incompressible State. " Thus, says he, if the
 " Particles of Water are turned into Vapour,
 " by repelling each other strongly, and repel
 " Air more than they repel each other; Ag-
 " gregates of such Particles, made up of Va-
 " pour and Vacuity, may rise in Air of different
 " Densities, according to their own Density
 " depending on their Degree of Heat." He
 observes farther, that Heat acts more power-
 fully on Water, than on common Air; for that
 the same Degree of Heat, which rarefies Air
 two Thirds, will rarefy Water near fourteen
 thousand Times, changing it into Steam or
 Vapour as it boils it. And in Winter, that
 small Degree of Heat, which in Respect of our
 Bodies appears cold, will raise a Steam or Va-
 pour from Water, at the same Time that it
 condenses Air. Lastly, he observes, That the
 Density and Rarity of this Vapour depends
 chiefly on its Degree of Heat, and but little
 on the Pressure of the circumambient Air. From
 all which he infers, That the Vapour being
 more rarefied near the Surface of the Earth,
 than the Air is there by the same Degree of
 Heat, must necessarily be buoyed up into the
 Atmosphere; and since it does not expand it-
 self much, though the Pressure of the incum-
 bent Air grows less, at length it finds a Place
 where the Atmosphere is of the same specific
 Gravity

Diff. VI *Ascent of Vapours, &c.* 135

Gravity with itself, and there floats, till by some Accident or other, it is converted again into Drops of Water, and falls down in Rain.

And to shew that Air is not necessary for the Formation of Steam or Vapour, he gives us this Experiment.

“ ABCD (*Fig. 37.*) is a pretty large Vessel of Water, which must be set upon the Fire to boil. In this Vessel must be suspended the Glass Bell E, made heavy enough to sink in Water ; but put in, in such a Manner, that it be filled with Water when upright, without any Bubbles of Air at its Crown within, the Crown being all under Water. As the Water boils, the Bell will by Degrees be emptied of its Water, being pressed down by the Steam, which rises above the Water in the Bell ; but, as that Steam has the Appearance of Air, in Order to know whether it be Air or not, take the Vessel off the Fire, and draw up the Bell by a String fastened to its Knob at Top, till only the Mouth remains under Water ; then as the Steam condenses by the cold Air on the Outside of the Bell, the Water will rise up into the Bell at F, quite to the Top, without any Bubble above it ; which shews, that the Steam, which kept out the Water, was not Air. ”

This

136 *Of the Formation and* Part II.

This Hypothesis, however plausible it may appear, is attended with Difficulties, as well as the other. For,

First, If the repulsive Power of the Particles of Water is sufficiently augmented by Heat as such (and that by a very small Degree of it) to make them recede from each other, and become specifically lighter than common Air; how comes it to pass, that all the Particles of Water, as soon as, or before it boils, have not their repulsive Forces thus augmented, since they are all under a much greater Degree of Heat, than is necessary to raise Vapour?

Secondly, Allowing that they may rise off from the Surface of the Water, and float in the circumambient Air, as being lighter than it, why do not their repulsive Forces, as they rise up into the Air, and the superincumbent Pressure becomes less, drive them to greater Distances from each other, and so cause them to continue lighter than the Air about them at all Heights?

Thirdly, If the Pressure of the Air about them, when near the Surface of the Earth, is not sufficient to bring them into so close Contact, as to form Drops of Water there, what Force will they find sufficient for it, when they are carried aloft into a Region of Air, where the Pressure is not near so great? The Doctor hints, that they are formed into Rain, "when
" by the great Diminution of the specific Gra-
" vity

Diff. VI. *Ascent of Vapours, &c.* 137

“ vity of the Air about a Cloud, it has a
“ great Way to fall, in which Case, he says,
“ the Resistance of the Air, which increases
“ as the Square of the Velocity of the de-
“ scending Cloud, causes the floating Particles
“ of Water to come within the Power of each
“ other’s Attraction, and form such big Drops
“ as, being specifically heavier than any Air,
“ must fall in Rain.” But as the inferior Air,
from the Cloud to a considerable Depth below
it, cannot be supposed to leave it all at once,
the Vapours must necessarily fall all the Way
through a resisting *Medium*; so that the little
Velocity the Cloud can acquire, while it is in
the Form of Vapour, will never produce a
Resistance from the Air greater than the Pres-
sure which the Vapours sustained below.

Lastly, The Experiment, brought to make
Way for this Hypothesis, shows clearly, that
Vapour formed without the Resistance of Air,
is immediately condensed into Water again, as
soon as it is suffered to cool: Which is widely
different from what happens to Vapours buoyed
up into the colder Regions of the Air, and
floating there, till they fall in Rain.

These are all, or the principal *Hypotheses*,
that have been framed for the Solution of this
common *Phænomenon*: Which as they seem
inadequate to the Effects produced, and there-
fore unsatisfactory, I thought it not amiss to
sug-

suggest to the Reader the chief Difficulties, with which I conceive them to be attended. But as it is easier to pull down, than to build up, to overturn a weak *Hypothesis*, than to raise and support a strong and sufficient one; so, I must own, in this Case, I can think of no Way of accounting for the Rise of Vapours, according to the received Principles of Philosophy, or say wherein their Nature consists. Upon this Account it is impossible I should give a Philosophical Account of their Resolution into Rain. It must suffice just to mention the Causes, which are observed to produce that Change.

The first is, That Part of the Air beneath them is carried away by contrary Winds blowing from the same Place*; for then the Remainder being too light to buoy them up, the upper ones, in all Probability, precipitate down upon the lower ones, unite with them, and forms Drops of Rain†. Another is, great
Quan-

* Perhaps it may be thought, that when the Air, which impends over any Place, is carried away from thence by contrary Winds, the Vapours which float in it should be carried away too; and so the few that remain should be easily supported; or, if they fall, should not produce much Rain.

Now, in Answer to this, it must be considered, that as the inferior Air is carried away from any Place by contrary Winds, the Height of the Atmosphere is diminished thereby; wherefore, in all Probability, the superior Air rushes in by a contrary Current to fill up the Vacuity at the Top, which bringing in its Contents of Vapour affords a continual Supply for the Rain that falls.

† A remarkable Instance we have of this, in that Part of the *Atlantic Ocean*, which the Sailors call the *Rains*, (see *Dissert. V.*)
from

Diff. VI. *Ascent of Vapours, &c.* 139

Quantities of them, being driven by the Winds against the Sides of Mountains, are thereby made to coalesce, and so constitute Rain*.

It

from the frequent Rains that fall there: the Occasion of which is; that the Atmosphere is diminished by continual contrary Winds blowing from thence, so that it is not able to sustain the Vapour it receives.

* Upon this depends the Difference of the Seasons of the Year at *Malabar* and *Coromandel* in the *East-Indies*, and at *Jamaica* in the West. See Dr. Gordon's Discourse on the Causes of the Change of Weather, Philosophical Transactions, N^o. 17. —

“ The Rivers of *Indus* and *Ganges*, says he, where they enter the Ocean, contain between them a large *Chersonesus*, which is divided in the Middle by a Ridge of Hills, which they call the Gate, which run along from East to West, and quite through to Cape *Coineri*. On the one Side is *Malabar*, and on the other *Coromandel*. On the *Malabar* Side, between that Ridge of Mountains and the Sea, it is after their Appellation Summer from September till April; in which Time it is always a clear Sky, without once, or very little Raining. On the other Side the Hills, on the Coast of *Coromandel*, it is at the same Time their Winter, every Day and Night yielding Abundance of Rain, and from April to September it is, on the *Malabar* Side their Winter, and on the other Side their Summer: So that in little more than twenty Leagues Journey in some Places, as where they cross the Hills to St. *Thomas*, on the one Side of the Hill you ascend with a fair Summer; on the other you descend with a stormy Winter. The like is said to be at Cape *Razalgate* in *Arabia*. And Dr. *Tropham* relates the same of *Jamaica*, intimating that there is a Ridge of Hills which runs from East to West, through the midst of the Island, and that the Plantations on the South Side of the Hills have, from November to April, a continual Summer, whilst those on the North Side have as constant a Winter, and the contrary from April to November.

“ From these and such like Accounts it seems evident, that a bare lessening of the Atmosphere's Gravity will not occasion Rain, but that there is also needful either a sudden Change of Winds, or a Ridge of Hills to meet the Current of the Air and Vapours, whereby the Particles of the Vapours are driven together, and so fall down in Drops of Rain. And hence it is,

T

“ that

It is generally thought, That if the Vapours in their Descent pass through a Region of Air sufficiently cold, they are there frozen into Icicles,

“ that whilst the Wind blows from the North-east, viz. from November to April (see the Account of the Monsoons in the foregoing Dissertation) “ there are continual Rains in the northerly Plantations of *Jamaica*, and on the Side of *Coromandel* in the *East-Indies*, because the Winds beat against that Side of the Hills; and so there is fair Weather on the other Side of these Hills, in *Malabar* and in the southern Plantations of *Jamaica*, there being no Winds to drive the Vapours together. “ But in the southerly Monsoons, viz. from April to November, “ *Malabar* and the southerly Plantations of *Jamaica* have Floods of Rain, the Wind beating against that Side of the Hills, whilst in *Coromandel*, and the other Side of *Jamaica*, there is fair and clear Weather.

“ This serves also to clear the Singularity of Seasons in *Peru*, beyond any other Parts of the Earth, and seems to be assigned by *Aristotle* as the Cause of it. *Peru* runs along from the Line Southwards about a thousand Leagues. It is said to be divided into three Parts, long and narrow, which they call *Lanos*, *Sierras*, and *Andes*; the *Lanos*, or Plains, run along the South-Sea Coast; the *Sierras* are all Hills with some Vallies; and the *Andes* steep and craggy Mountains. The *Lanos* have some ten Leagues in Breadth, in some Parts less, and in some more; the *Sierras* contain some twenty Leagues in Breadth, the *Andes* as much, sometimes more, sometimes less; they run in Length from North to South, and in Breadth from East to West. This Part of the World is said to have these remarkable Things: 1. All along the Coast, in the *Lanos*, it blows continually with one only Wind, which is South and South-west, contrary to that which usually blows under the torrid Zone. 2. It never rains, thunders, snows, or hails in all this Coast, or *Lanos*, though there falls sometimes a small Dew. 3. Upon the *Andes* it rains almost continually, though it be sometimes more clear than other. 4. In the *Sierras*, which lie between both Extrems, it rains from September to April, but in the other Seasons it is more clear, which is when the Sun is farthest off, and the contrary when it is nearest. Now the Reason of all seems to be this; The eastern Breezes, which blow constantly under the Line, being stopp'd in their

“ Court

Diff. VI. *Ascent of Vapours, &c.* 141

Icicles, and from Snow. But this Opinion seems to be false; because it frequently snows when the Barometer is high, at which Time the Vapours cannot begin to fall. It is therefore more probable, that they are first frozen into Icicles, and by that means shooting forth into several Points every Way from the Center (agreeably to the Nature of Freezing) lose their Form; and so becoming specifically heavier than Air fall down, and in their Passage, several being congealed together, form Fleeces of Snow*.

Hail is evidently no other, than Drops of Rain congealed into Ice. This happens, when in their Passage through the inferior Air, they

“ Course by the *Sierras* and *Andes*, and yet the same Breezes being to be found in the South Sea beyond *Peru*, as appears by the easy Voyages from *Peru* to the *Philippines*, a Current of Wind blows from the South on the Plains of *Peru*, to supply the eastern Breeze on the South-Seas, and there being but one constant Gale on these Plains, and no contrary Winds or Hills for it to beat upon, this seems to be the Reason why the Vapours are never or very seldom driven into Rain. And the *Andes* being as high perhaps in many Places as the Vapours ascend in the highest Degree of the Atmosphere's Gravity, this may probably be the Reason, why the eastern Breeze, beating constantly against these Hills, occasions Rain upon them at all Seasons of the Year. And the *Sierras* being it seems lower than the *Andes*, therefore from *September* to *April*, when the Sun is nearest, and so the Atmosphere's Gravity less, and the Vapours lower, they are driven against the *Sierras* into Rain.

The like is to be said of other Countries. They are all, *cæteris paribus*, more or less rainy, as they are more or less mountainous.

Egypt, which is quite without Mountains, has seldom or never any Rain; but is plentifully watered by the *Nile*, which yearly rises above its Banks, and overflows the whole Country.

* See a Discourse on the Nature of Snow, Philosophical Transactions N°. 92.

meet with *nitrous* Particles, which are known to contribute greatly to Freezing. Their Magnitude is owing to a fresh Accession of Matter, as they pass along. Hence we see the Reason why Hail is so frequent in Summer, because at that Time greater Quantities of Nitre are exhaled from the Earth, and float up and down in the Air.

If the Vapours, after they are exhaled from off the Waters, do not rise very high in the Atmosphere, but hover near the Surface of the Earth, they then constitute what we call a Fog. And, if they ascend higher, they still appear in the same Form to those, who, being upon the Tops or Sides of Mountains, are at an equal Height with them; though to those, who are below, they appear as Clouds.

If they fall to the Earth, without uniting into Drops large enough to be called Rain, they are then said to fall in Dew.

See farther on this Subject, *Derham's* *Physi. Theolog.* Book I. Chap. 3, and Book II. Chap. 5. *Spectacle de la Nature*, Dialog. 21, and 23. *Nieuwentyt* *Contempl.* 19. *Clare's* *Motion of Fluids.* *Regnault*, Vol. III. *Conversat.* 7. *Musschenbroek* *Epitom. Phys. Mathemat.* Par. II. Cap. 24. *Melchior Verdries* *Physic. Pars special.* Cap. V. §. 8. And the Authors referred to in Mr. *Johnson's* *Questiones Philosoph.* Cap. IV. Quæst. 7.

DISSERTATION VII.

Of the Causes of Thunder and Lightning, and of the Aurora Borealis.

THOSE Philosophers, who maintain that Vapours are buoyed up into the Air, by Particles of Fire adhering to them (as explained in the foregoing Dissertation) account for the *Phænomena* of Thunder and Lightning, in the following Manner. They suppose, that from the Particles of Sulphur, Nitre, and other combustible Matter, which are exhaled from the Earth, and carried into the higher Regions of the Atmosphere, together with the ascending Vapours, is formed an inflammable Substance ; which, when a sufficient Quantity of fiery Particles is separated from the Vapour by the Collision of two Clouds, or otherwise, takes Fire, and shoots out into a Train of Light, greater or less, according to the Strength and Quantity of the Materials.

This Opinion is false for the Reasons mentioned in the foregoing Dissertation, which plainly show, that it is impossible the Vapours should be attended with such fiery Particles, as is here supposed.

Neither

144 *Of the Causes of Thunder* Part II.

Neither have we Occasion to fly to such an *Hypothesis*; for, as Vapours exhaled from the Surface of the Water are carried up into the Atmosphere: in like Manner, the *Effluvia* of solid Bodies are continually ascending thither: Now we find by Experiment, that there are several inflammable Bodies, which, being mixed together in due Proportion, will kindle into Flame by Fermentation alone,* without the Help of any fiery Particles †. When therefore

* See the Theory of Fermentation explained in the following Dissertation.

† Monsieur *Lemery* having covered up in the Earth about fifty Pounds of a Mixture composed of equal Parts of Sulphur, and Filings of Iron tempered with Water; after eight or nine Hours Time, the Earth, where it was laid, vomited up Flames. *Hist. de l'Acad.* 1700, p. 574.

From this Experiment we see the true Cause of the Fire of *Etna* and *Vesuvius*, and other burning Mountains. They probably are Mountains of Sulphur, and some other Matter proper to ferment with it, and take Fire. From like Causes proceeds the Heat of Bath-waters, and other hot Springs.

Mix a small Quantity of Gun-Powder with Oyl of Cloves, pour gently upon this Mixture, two or three times as much Spirit of Nitre, and you will observe a bright Inflammation suddenly arising from it. A Mixture of the two Fluids alone will take Fire; the Powder is added only to augment the Inflammation.

Take two Pounds of Nitre, or refined Salt Petre well dried and reduced to the finest Powder, with a Pound of Oyl of common Vitriol: From this Mixture may be drawn by Distillation a Spirit of Nitre capable of inflaming Oyl of Turpentine. *Mem. de l'Acad.* 1726, p. 97, &c. Put into a Glass an Ounce of this Spirit of Nitre, with an Ounce of Oyl of Vitriol; pour upon it an equal Quantity of Oyl of Turpentine, and a very fine Flame will arise suddenly with a great Explosion. When the Liquors are fresh the Effect is much greater. If we mix a Dram of the Spirit of Nitre and three of Oyl of Turpentine, with only one of the Spirit

Diff. VII. and Lightning, &c. 145

fore there happens to be a proper Mixture of the *Effluvia* of such Bodies floating in the Air, they ferment, kindle, and flashing like Gunpowder, occasion those Explosions, and Streams of Fire, which we call Thunder and Lightning.

As to the particular Species of *Effluvia*, which compose this Mixture, that cannot exactly be determined; they are thought to be chiefly sulphureous and nitrous: *sulphureous*, because of the sulphureous Smell which Lightning generally leaves behind it, and of that sultry Heat in the Air which is commonly the Fore-runner of it: *nitrous*, because we don't know of any Body so liable to a sudden and violent Explosion, as Nitre is *.

The

of Vitriol, the Mixture will take Fire immediately. If the same Experiment be made with the Balm of *Mecca*, a sudden Flame will arise, with a Noise like that of the Report of a Gun.

There are two celebrated Experiments of this Kind, though they do not come up exactly to the present Purpose, because they will not succeed, unless the Ingredients be first heated, the one of *Aurum fulminans*, and the other of *Pulvis fulminans*. The first is a Mixture of Salt of Tartar, and Gold dissolved by *Aqua Regia*. A small Quantity of this, if put into a Brass Spoon, and heated with the Flame of a Candle, causes a sudden Noise resembling that of Thunder; and goes off with great Violence. The other is a Mixture of three Parts of Nitre, two of Salt of Tartar, and one of Sulphur. This Mixture when set upon the Fire, and warmed to a certain Degree, is dissipated all on a sudden with great Thundring, like the *Aurum fulminans*.

See an Account of Exhalations taking Fire of their own Accord in Cole-Pits. *Power's Experimental Philosophy*, p. 62. and 181.

* Dr. *Lifter* is of Opinion, That the Matter both of Thunder and Lightning, and also of Earthquakes, is the *Effluvia* of the *Pyrites*,

146 *Of the Causes of Thunder.* Part II

The Effects of Thunder and Lightning are owing to the sudden and violent Agitation the Air is put into thereby, together with the Force of the Explosion *; and not to Thunderbolts falling from the Clouds; as supposed by the Vulgar †.

Pyrites; as he does, that the Matter of *Vulcano's* is the *Pyrites* itself. This is a Mineral that emits copious Exhalations, and is exceedingly apt to take Fire upon the Admission of Moisture. See the Doctor's Defence of his Notion in the Philosophical Transactions, N^o 157. He thinks this may be the Reason why *England* is so little troubled with Earthquakes, and *Italy*, and almost all Places round the *Mediterranean Sea*, so very much, *viz.* because the *Pyrites* are rarely found in *England*; and where they are, they lie very thin, in Comparison of what they do in those Countries; as the vast Quantity of Sulphur, emitted from the burning Mountains there, seems to shew.

* Lightning is said to have dissolved Silver, without burning the Purse it was in; and to have melted the Sword, without touching the Scabbard, and the like. The Occasion of this may possibly be, that the Matter of the Exhalation may be so subtle and penetrating, that, as we see it happens with *Aqua Fortis*, or *volatile Salts*, it may pass through soft Bodies without altering their Texture, while it spends its whole Force on hard ones, in which it finds the greater Resistance.

† Some are inclined to think, that Thunderbolts are artificial, and that they were applied by the Ancients to some Use. What confirms them in their Opinion, is, that they are found more frequently where Sepulchres have been, than in other Places. And, as it was the Custom of the Ancients to have their Arms buried with their Ashes, they think they might be of some Use in War. Some are of Opinion, they were used in Sacrifices. See *Melchior Verdrie's* Physic. Pars special. Cap. V. §. 9. *Hedelius* Exercit. Med. Philol. Cont. II. Dec. I. p. 103. *Schminckius* Profess. Marburg. Dissertat. de Urnis Sepulchralibus, & Armis Lapideis, A. 1714. *Herman Nunningius* Sepulchret. Westphal. Mimigard. Gentil. p. 44. *Jo. Henr. Coharsen* Ollieg. Hiltor. Physic. p 44.

Diff. 7. *Of the Aurora Borealis.* 147

The Distance the Thunder is from us, may nearly be estimated by the Interval of Time between our seeing the Lightning, and hearing the Thunder. For, as the Motion of Light is so very quick, that the Time it takes up, in coming to us from the Cloud, is not perceptible; and as that of Sound is about a thousand Feet in a Second; allowing a thousand Feet for every Second, that passes between our seeing the one, and hearing the other; we have the Distance of the Cloud, pretty nearly, from whence the Thunder comes.

We sometimes see Flashes of Lightning, though the Sky be clear and free from Clouds; in this Case they proceed from Clouds, that lie immediately below our Horizon.

Of Affinity with the *Phænomena* of Lightning are those of the *Aurora Borealis*, or *Northern Lights*, which, of late Years, have very frequently appeared in our Climate*. These Lights differ so much from each other, that to give a Description of one alone, would not be

* *Phænomena* of this Kind are reported to have been very frequent in *Groenland*, *Iceland*, and *Norway*, Countries near the Pole. The only ones, that are upon Record, as having appeared in *England*, before that of *March* the 6th, 17 $\frac{1}{2}$, are those of *January* the 30th, 1560, *October* the 7th, 1564. *November* 14th and 15th, 1574, and a small one seen near *London* on the 9th of *August*, 1708. On *November* the 16th, 1707, a small one appeared in *Ireland*. Since that of *March* the 6th, 17 $\frac{1}{2}$, they have been, and still continue very common.

148 *Of the Aurora Borealis. Part II.*

sufficient to acquaint the Reader with all the Circumstances wherewith they are attended. I shall therefore collect together such *Phænomena*, as have been most generally observed, and reduce them to the ten following Propositions, adding in the Notes, by Way of *Specimen*, a full Account of that most remarkable *Aurora*, which was seen *March* the 6th 17¹³/₁₆, as it was laid before the *Royal Society* by Dr. *Halley*, at their Request *.

The

* “ On *Tuesday* the 6th of *March*, in the Year 1716 (the Afternoon having been very serene and calm, and somewhat warmer than ordinary) about the Time it began to grow dark (much about seven of the Clock) not only in *London*, but in all Parts of *England*, where the Beginning of this wonderful Sight was seen; out of what seemed a dusky Cloud, in the North-east Parts of the Horizon, and scarce ten Degrees high, the Edges whereof were tinged with a reddish Yellow, like as if the Moon had been hid behind it, there arose very long luminous Rays, or Streaks perpendicular to the Horizon, some of which seemed nearly to ascend to the *Zenith*. Presently after, that reddish Cloud was swiftly propagated along the northern Horizon into the North-west, and still farther westerly; and immediately sent forth its Rays from all Parts, now here, now there, they observing no Rule or Order in their rising. Many of those Rays, seeming to concur near the *Zenith*, formed there a *Corona*, or Image, which drew the Attention of all Spectators. Some likened it to that Representation of Glory, wherewith our Painters in Churches surround the Holy Name of God. Others to those radiating Stars, wherewith the Breasts of *Knights* of the Order of the *Garter* are adorned. Many compared it to the *Concave* of the great *Cupola* of *St. Paul's* Church, distinguished with Streaks alternately light and obscure, and having in the Middle a Space less bright than the rest, resembling the *Lanthorn*. Whilst others, to express as well the Motion as Figure thereof, would have it to be like the Flame in an Oven, reverberated and rolling against the arches

“ Rai

Diff. VII. *Of the Aurora Borealis.* 149

The most general *Phænomena* of an *Aurora Borealis* are these that follow.

I. In the northern Parts of the *Horizon*, there is commonly the Appearance of a very black

“ Roof thereof: Some thought it liker to that tremulous Light,
“ which is cast against the Cieling by the Beams of the Sun, re-
“ flected from the Surface of the Water in a Bason, that's a little
“ shaken. But all agree, that this *Spectrum* lasted only a few
“ Minutes, and exhibited itself variously tinged with Colours, Yel-
“ low, Red, and a dusky Green: Nor did it keep in in the same
“ Place; for when first it began, it appeared a little to the
“ Northwards of the *Zenith*, but by Degrees declining towards
“ the South, the long *Striae* of Light, which arose from all Parts
“ of the northern Semicircle of the *Horizon*, seemed to meet to-
“ gether, not much above the Head of *Castor*, or the northern
“ *Twin*, and there soon disappeared.

“ After the first *Impetus* of the ascending Vapour was over, the
“ *Corona* appeared no more; but still, without any Order, as to
“ Time or Place, or Size, luminous *Radii*, like the former, con-
“ tinued to arise perpendicularly, now oftener, and again sel-
“ domer; now here, now there; now larger, now shorter.
“ Nor did they proceed as at first, out of a Cloud, but oftener
“ would emerge at once out of the pure Sky, which was more
“ than ordinary serene and still. Nor were they all of the same
“ Form. Most of them seemed to end in a Point upwards, like
“ erect Cones; others like truncate Cones, or Cylinders, so much
“ resembling the long Tails of Comets, that at first Sight, they
“ might well be taken for such. Some of those Rays would con-
“ tinue visible for several Minutes; when others, and those the
“ much greater Part, just shewed themselves, and died away.
“ Some seemed to have little Motion, and to stand, as it were,
“ fixed among the Stars, whilst others, with a very perceptible
“ Translation, moved from East to West under the Pole, contrary
“ to the Motion of the Heavens; by which Means they would
“ sometimes seem to run together, and at other Times to fly one
“ another.

“ After this Sight had continued about an Hour and a half,
“ those Beams began to rise much fewer in Number, and not near
“ so high; and by Degrees, that diffused Light, which had illus-

150 *Of the Aurora Borealis.* Part II.

black Cloud; but it is evident that it is no real Cloud, because the smallest Stars are visible through it. This apparent Cloud is extended sometimes farther towards the West, than

“trated the northern Parts of the Hemisphere, seemed to subside,
 “and settling on the Horizon, formed the Resemblance of a very
 “bright *Crepusculum*. That this was the State of this *Phænomenon*,
 “in the first Hours, is abundantly confirmed by the unanimous
 “Consent of several. For, by the Letters we have received from
 “almost all the extreme Parts of the Kingdom, there is found very
 “little Difference from what appeared at *London* and *Oxford*; un-
 “less that in the North of *England*, and in *Scotland*, the Light
 “seemed somewhat stronger and brighter.

“Hitherto I have related the Observations of others; as to
 “myself, I had no Notice of this Matter, till about nine of the
 “Clock: I immediately perceived toward the South and South-
 “west Quarter, that though the Sky was clear, yet it was tinged
 “with a strange Sort of Light; so that the smaller Stars were
 “scarce to be seen, and much as it is when the Moon of four
 “Days old appears after Twilight. I perceived at the same Time
 “a very thin Vapour to pass before us, which arose from the pre-
 “cise East Part of the Horizon, ascending obliquely, so as to
 “leave the *Zenith* about fifteen or twenty Degrees to the North-
 “ward. But the Swiftness, wherewith it proceeded, was scarce to
 “be believed, seeming not inferior to that of Lightning; and
 “exhibiting, as it passed on, a Sort of momentaneous *Nubecula*,
 “which discover’d itself by a very diluted and feint Whiteness;
 “and was no sooner formed, but before the Eye could well take
 “it, it was gone, and left no Signs behind it. Nor was this a
 “single Appearance; but for several Minutes, about six or seven
 “Times in a Minute, the same was again and again repeated;
 “these Waves of Vapour regularly succeeding one another, and
 “at Intervals very nearly equal; all of them in their Ascent pro-
 “ducing a like transient *Nubecula*.

“By this Particular we were first assured; that the Vapour we
 “saw, became conspicuous by its own proper Light, without the
 “Help of the Sun’s Beams; for these *Nubeculae* did not discover
 “themselves in any other Part of their Passage, but only between
 “the South-east and South, where being opposite to the Sun,
 “they

Diff. VII. *Of the Aurora Borealis.* 151

than to the East; sometimes farther towards the East, then to the West; and generally takes up a Quarter of the Horizon, more or less.

2. The

“ they were deepest immerfed in the Cone of the Earth’s Shadow;
“ nor were they vifible before or after. Whereas the contrary
“ muft have happened, had they borrowed their Light from the
“ Sun.

“ On the western Side of the northern Horizon, *viz.* between
“ West and North-west, not much past ten of the Clock, I ob-
“ served the Representation of a very bright Twilight, contiguous
“ to the Horizon, out of which arose very long Beams of Light,
“ not exactly erect towards the *Vertex*, but something declining
“ towards the South; which, ascending by a quick and undulating
“ Motion to a confiderable Height, vanished in a little Time;
“ whilst others, though at uncertain Intervals, fupplied their
“ Place. But at the fame Time, through all the reft of the nor-
“ thern Horizon, *viz.* from the North-west to the true East,
“ there did not appear any Sign of Light to arife from, or join
“ to, the *Horizon*; but what appeared to be an exceeding black
“ and difmal Cloud, feemed to hang over all that Part of it. Yet
“ was it no Cloud, but only the ferene Sky, more than ordinary
“ pure and limpid, fo that the bright Stars fhone clearly in it,
“ and particularly *Canuda Cygni*, then very low in the North;
“ the great Blacknefs manifeltly proceeding from the Neighbour-
“ hood of the Light, which was collected above it. For the
“ Light had now put on a Form quite different from all that we
“ have been describing, and had fashioned itfelf into the Shape of
“ two *Laminae*, or Streaks, lying in a Pofition parallel to the *Hori-*
“ *zon*, whose Edges were but ill-terminated. They extended them-
“ felves from the North by East to the North-east, and were each
“ about a Degree broad; the undermoft about eight or nine Degrees
“ high, and the other about four or five Degrees over it; thefe
“ kept their Places for a long Time, and made the Sky fo light,
“ that I believe a Man might eafily have read an ordinary Print
“ by the Help thereof.

“ Whilst I was viewing this furprizing Light, and expecting
“ what was farther to come, the northern End of the upper *La-*
“ *mina* by degrees bent downwards, and at length clofed with
“ the End of the other that was under it, fo as to fhut up on the
“ North-Side an intermediate Space, which ftill continued open
“ to

152 *Of the Aurora Borealis. Part II.*

2. The upper Edge of this Cloud (which is sometimes within less than fix Degrees of the Horizon, and sometimes forty or fifty above it)

“ to the East. Not long after this, in the said included Space, I
 “ saw a great Number of small Columns, or whitish Streaks, to
 “ appear suddenly erect to the Horizon, and reaching from the
 “ one *Lamina* to the other; which instantly disappearing, were
 “ too quick for the Eye, so that I could not judge, whether they
 “ arose from the under, or fell from the upper; by their sudden
 “ Alterations, they made such an Appearance, as might well
 “ enough be taken to resemble the Conflict of Men in Battle.

“ And much about the same Time, there began on a sudden to
 “ appear, low under the Pole, and very near due North, three or
 “ four lucid *Areas*, like Clouds, discovering themselves in the
 “ pure but very black Sky, by their yellowish Light. These, as
 “ they broke out at once, so after they had continued a few Mi-
 “ nutes, disappeared as quick, as if a Curtain had been drawn
 “ over them: Nor were they of any determined Figure, but
 “ both in Shape and Size might properly be compared to small
 “ Clouds illuminated by the full Moon, but brighter.

“ Not long after this, from above the foresaid two *Laminae*,
 “ there arose a very great *Pyramidal* Figure, like a Spear, sharp
 “ at the Top, whose Sides were inclined to each other, with an
 “ Angle of about four or five Degrees, and which seemed to reach
 “ up to the *Zenith*, or beyond it. This was carried with an
 “ equable, and not very slow Motion, from the North-east where
 “ it arose, into the North-west, where it disappeared, still keep-
 “ ing in a perpendicular Situation, or very near it; and passing
 “ successively over all the Stars of the *little Bear*, did not efface
 “ the smaller ones in the Tail, which are of the fifth Magni-
 “ tude; such was the extream Rarity, and Perspicuity of the
 “ Matter whereof it consisted.

“ This single Beam was very remarkable for its Height above
 “ all those, that, for a great while before, had preceeded it, or
 “ that followed it.

“ It being now past eleven of the Clock, and nothing new of-
 “ fering itself to our View, but repeated *Phases* of the same Spec-
 “ tacle; I observed, that the two *Laminae*, or Streaks, parallel
 “ to the *Horizon*, had now wholly disappeared; and the whole
 “ Spectacle reduced itself to the Resemblance of a very bright
 “ Cre-

Diff.VII. Of the Aurora Borealis. 153

it) is generally terminated with a very lucid Arch, from one to four or five Degrees broad, whose Center is below the Horizon. Sometimes

“ *Crepusculum* setting on the Northern *Horizon*, so as to be
 “ brightest and highest under the Pole itself; from whence it
 “ spread both Ways into the North-east and North-west. Under this, in the Middle thereof, there appeared a very black
 “ Space, as it were the Segment of a lesser Circle of the Sphere
 “ cut off by the *Horizon*. It seemed to the Eye like a dark Cloud,
 “ but was not so; for by the Telescope the small Stars appeared
 “ through it more clearly than usual, considering how low they
 “ were: And upon this as a *Basis*, our *Lumen Auroriforme* rested,
 “ which was no other than a Segment of a Ring, or *Zone*
 “ of the Sphere, intercepted between two parallel lesser Circles,
 “ cut off likewise by the *Horizon*; or the Segment of a very
 “ broad *Iris*, but of one uniform Colour, *viz.* a Flame-Colour
 “ inclining to Yellow, the Center thereof being about forty Degrees below the *Horizon*. And above this there were seen
 “ some Rudiments of a much larger Segment, with an Interval
 “ of dark Sky between, but this was so exceeding feint and
 “ uncertain, that I could make no proper Estimate thereof.

“ I attended this *Phænomenon* till near three in the Morning, and
 “ the Rising of the Moon: But for above two Hours together it
 “ had no Manner of Change in its Appearance, nor Diminution,
 “ nor Increase of Light; only sometimes, for very short Intervals,
 “ as if new Fewel had been cast on a Fire, the Light seemed to
 “ undulate and sparkle not unlike the rising of a vaporous Smoak
 “ out of a great Blaze when agitated. But one Thing I assured
 “ myself of, that the *Iris*-like Figure did by no means owe its
 “ Origin to the Sun’s Beams: For that about three in the Morning,
 “ the Sun being in the Middle between the North and East,
 “ our *Aurora* had not followed him, but ended in that very Point
 “ where he then was: Whereas in the true North, which the
 “ Sun had long passed, the Light remained unchanged, and in
 “ its full Lustre.

Appearances of this Kind have been taken Notice of both by
Pliny, *Seneca* and *Aristotle*; the last of which calls the vibrating Light near the *Zenith*, *Δαλοι*; the more steady perpendicular Streams, *Δοκός*; and the *Aurora* itself, from the apparent dark Cloud just below it, *Χάσμα*. That *Aurora* which was observed
 by

154 *Of the Aurora Borealis. Part II.*

times there are two or more of these Arches, one above another. In some, the Cloud is not terminated by an Arch, but by a long bright Streak

by Monsieur *Gassendi* in *Provence*, on the 21st of *September*, in the Year 1621, was very remarkable, at that Time.

He tells us, That about the End of Twilight in the Evening, when the Sky was very clear, and there was no Moon, there appeared in the North a Sort of a rising Morn, which ascending by Degrees, became intermingled with certain Streaks, as it were, or Rays perpendicular to the Horizon: And that at the same Time there appeared some small *passing* whitish Clouds between the South and the Place of the Sun's setting in Winter; and that in the Place where the Sun sets in Summer, a bright Redness seemed to arise in the Form of a *Pyramid*, which moved towards the setting of the Sun at the *Equinox*; and which was distinguished into three several *Pyramids*, which in a little Time were confounded one with another, and at last disappeared. When this Redness ceased, the northern Whiteness was risen forty Degrees and more, that is, about the Altitude of the Pole-Star, forming itself into an Arch, and taking up near sixty Degrees of the *Horizon*. After this, certain *Chevrons*, or Columns of Rays, some whiter, and some a little darker, began more plainly to be distinguished, of about two Degrees in Breadth, and perpendicularly posited; so that all that Part appeared as it were fluted. The whole Circumference immediately appeared scalloped; and then some of those Columns which were in the Middle, and that were the whitest, began as it were to leave their Places with great Impetuosity, and in less than a Quarter of a Minute, raised themselves almost to the Top, putting on the Form of *Pyramids*, which they would retain four or five Minutes. It was about nine of the Clock, when the Arch of Whiteness began to decrease or sink; at which Time certain very white Streams of Smoke began to issue out from the Columns which were remaining under the *Pyramids*, and darting upwards with very great Rapidity through the *Pyramids*, like Javelins, vanished immediately when they came to the Tops of them. This Spectacle lasted about an Hour; after which the Whiteness sunk down to about six Degrees of the *Horizon*. *Vide Abregé de Gassendi*, Tom. V. P. 245.

This

Diff. VII. Of the Aurora Borealis. 155

Streak of Light, lying parallel to the *Horizon*. The Limb of this luminous Arch, or parallel Streak is not always even and regular, but sinks lower in some Parts, than in others.

3. Out of this Arch proceed Streams of Light, generally perpendicular to the *Horizon*, but sometimes a little inclined to it. Most of them seem to end in a Point, like *Pyramids* or *Cones*; and often very much resemble the Tails of *Comets*. Sometimes there is no luminous Arch, nor Streak of Light; and then the Streams seem to issue out from behind the dark Cloud, being distinct from each other at their *Bases*.

4. The upper Ends of the Streams incessantly appear and vanish again, as quick as if a Curtain were drawn backwards and forwards before them; which sometimes causes such a seeming trembling in the Air, that you would

This *Phænomenon* appeared not only to *Gassendi* in *Provence*, but was seen at Places very distant from thence, as at *Tolose*, *Montaubon*, *Bordeaux*, *Grenoble*, *Dijon*, *Paris*, and *Rodn*, &c. and at most other Places in *France*, and the neighbouring Countries, that lie to the Northwards of *Provence*, unless where it was intercepted by Clouds; but no where in such as lie at any great Distance to the Southwards of it.

Monsieur *Gassendi* is thought to have given the Name of *Aurora Borealis* to this *Phænomenon*; but this is observed by Monsieur *Mairan*, to be a Mistake. See Mr. *Mairan*'s *Physical and Historical Treatise of the Aurora Borealis*, in the *Memoires de l'Academie Royale des Sciences*, Année 1731. or an Abstract of it in *Philosoph. Transact.* N^o. 431.

156 *Of the Aurora Borealis.* Part II.

imagine the upper Part of the Heavens to be, as it were, in Convulsions*.

5. They sometimes seem to meet in the *Zenith*, or more commonly to the Southward of it, about ten Degrees, more or less (sometimes they deviate a little to the South-east of the *Meridian*, and sometimes to the South-west); and there curling round, in some Measure, imitate Flame confined under an Arch; and being frequently tinged with various Orders of Colours, exhibit a most beautiful Appearance, resembling a Canopy finely painted†, of about ten or twenty Degrees in Breadth.

In many *Aurora*'s, there are Streams ascending from those Parts of the Heavens, which lie several Degrees to the Southwards of the Canopy; and in some, they appear to arise, though very rarely, almost as large, and numerous from the southern, as from the northern Parts of the Horizon.

6. The Height of the *Aurora Borealis* is very great; for that of *March* the sixth 17¹⁸ was visible from the West Side of *Ireland*, to the Confines of *Russia* and *Poland* on the East, and perhaps farther; extending at least over thirty Degrees of Longitude, and in La-

* See their Motions well described in the Account we have of an *Aurora* in the Philosoph. Transact. N^o. 395, Art. 2.

† See the various Colours of the Canopy, as well as those of an *Aurora* itself, accurately described by Pi. Greenwood, in Philosoph. Transact. N^o. 418, Art. 1.

Diff. VI. *Of the Aurora Borealis.* 157

titude from about the fiftieth Degree over almost all the North of *Europe*, and at all Places, exhibiting, nearly at the same Time, the same Appearances.

7. These Appearances have always been very frequent (as far as the Accounts we have of them inform us) in Countries that lie in, or near the *frigid Zone*, but very rare in those of our Latitude; they are now become very frequent with us, but always seem to proceed from the North; and are as yet unknown to the Inhabitants of the more southern Parts of our *Hemisphere*. Whether they are seen to those, who inhabit in, or near the other *frigid Zone*, is to us unknown.

8 In some, there are Trains of Light running horizontally, sometimes from the Middle to the Extremes, and sometimes from one Extreme to the other. And from these Trains often arise Streams perpendicular to the *Horizon*, and accompanying them as they pass along,

9. When all the Streaming is over, the *Aurora Borealis* commonly degenerate into a bright Twilight in the North, and there gradually dies away.

10. It is probable, that these *Phænomena* often happen in cloudy Nights, though we are not sensible of them; for 'tis observable, that in such Nights, there is frequently more Light

158 *Of the Aurora Borealis.* Part II.
than what usually proceeds from the Stars
alone.

The most obvious Solution of the *Aurora Borealis*, or at least what would appear so, to such as have only attended to the Circumstances of some particular ones, and which has accordingly been affixed by several * to their Accounts of the *Aurora's* they have seen, is that it is a thin Nitro-sulphureous Vapour, raised in our Atmosphere considerably higher than the Clouds; that this Vapour by Fermentation takes Fire, and the Explosion of one Portion of it kindling the next, the Flashes succeed one another, till the whole Quantity of Vapour within their Reach is set on Fire.

* Professor Cotes, at the End of his Description of a *Phænomenon* of this Kind, inserted in the Philosophical Transactions N^o. 365, observes, that supposing a Fund of Vapours or Exhalations at a considerable Height above us to be diffused every Way into a large and spacious Plane, parallel to the *Horizon*, that Fund of mixed Matter by Fermentation will emit Streams; and that if the Wind be still, they will ascend perpendicularly upwards; otherwise they will be inclined towards that Point of the *Horizon* which is opposite to that from which the Wind blows; and that they will appear, by the Rules of Perspective, in the first Case, to converge to the Spectator's *Zenith*, in the other, to some Point not far from it; and that if this Fund of Vapours be situated more to the North than the South, it will produce Streams of Light attended with such Circumstances, as then appeared: But he does not say, why the Vapours should be situated rather to the North than the South, or proceed to account for all the *Phænomena* of the *Aurora Borealis* in general from thence.

Some

Diff. VII. *Of the Aurora Borealis.* 159

Some have thought, that Vapours rarefied exceedingly by subterraneous Fire, and tinged with sulphureous Steams, might from thence be disposed to shine in the Night, and rising up to the Top of the Atmosphere, or even beyond its Limits (as we find the Vapours of Gun-powder, when heated in *Vacuo*, will shine in the Dark, and ascend to the Top of the Receiver, though exhausted) might produce those Undulations in the Air, which constitute this *Phænomenon*.

But these *Hypotheses* have been rejected, as insufficient: it having been thought impossible to account for all the Circumstances of the *Aurora* by them *.

* In the Commentaries of the Academy of Sciences at *Petersburgh*, I find a late Solution of the *Aurora Borealis* from Exhalations fermenting and taking Fire in the Atmosphere, which the Author *Chr. Maier* says, occasion the Appearance of the lucid Arch in the North, and thinks that the Streams, which seem to issue from thence, are no other than the Light of that Arch reflected to us from the under Side of some thin Clouds, that lie above it. As to its appearing in the North rather than in the South, he supposes that may be owing to the Cold of those Regions condensing the Exhalations, and thereby rendering them more liable to ferment than they are in the southern; but acknowledges ingenuously, that he has no Reason to suppose this, but its being necessary to his Solution. At the End he tells, That it was known in ancient Times as well as lately: But omits taking Notice, that it appears much oftener of late Years in our Climate than it used to do; and so avoids accounting for that Particular. *Vide Commentar. Academ. Scientiar. Imperial. Petropolitan. Tom. I. p. 351.*

Dr. *Halley* therefore has Recourse to the *magnetic Effluvia* of the Earth, which he supposes to perform the same Kind of Circulation with Regard to it, as the *Effluvia* of any particular *Terrella* * do with respect to that, *viz.* that they enter the Earth near the South Pole, and pervading its Pores, pass out again at the same Distance from the northern : And thinks, they may sometimes, by the Concourse of several Causes very rarely coincident, and to us as yet unknown, be capable of producing a small Degree of Light, either from the greater Density of the Matter, or perhaps from the greater Velocity of its Motion ; after the same Manner, as we see the *Effluvia* of *Electric* Bodies emit Light in the Dark.

Monsieur de *Mairan* has given us a Physical and Historical Treatise of the *Aurora Borealis*, and endeavours to prove that it is owing to the *Zodiacal Light*, or the Atmosphere of the Sun, spread on each Side of it along the *Zodiac* in the form of a *Pyramid*. This, he says, is sometimes extended to such a Length as to reach beyond the Orbit of our Earth, and then mixing itself with our Atmosphere, and being of an *Heterogeneous* Nature, produces

* A round Magnet, so called from the Resemblance it bears to the Earth.

Diff. VII. Of the Aurora Borealis. 161

the several Appearances, which are observed in the *Aurora Borealis* *.

I have just mentioned these two Solutions, because they come from two very ingenious Philosophers; though I doubt not but the Reader will agree with me, that they are much too fine spun to hold, and that they are no other than the ingenious *Reveries* of Persons determined to frame an Hypothesis at any Rate. I question not, but we may find Materials for the *Aurora Borealis*, without going so far for them, as these Gentlemen have done; and in particular that we have no Occasion to have Recourse either to the *magnetic Effluvia* of the Earth, or the *Zodiacal Light*, the Nature of both which we are wholly unacquainted with. The Materials employed in the first Solution (I mean such *Effluvia* as are continually exhaled from the Surface and Bowels of the Earth) if rightly considered, will afford a very easy and natural one, as I shall endeavour to shew in the following Manner.

First, we are assured by Experiment, that there are some Steams (as inflammable sulphureous ones) which are capable of so great a Degree of Expansion, that they will render themselves lighter than the Air they float in, though it

* See his Account at large, referred to at the End of Note, Page 155.

160 *Of the Aurora Borealis. Part II.*

Dr. Halley therefore has Recourse to the *magnetic Effluvia* of the Earth, which he supposes to perform the same Kind of Circulation with Regard to it, as the *Effluvia* of any particular *Terrella* * do with respect to that, *viz.* that they enter the Earth near the South Pole, and pervading its Pores, pass out again at the same Distance from the northern: And thinks, they may sometimes, by the Concourse of several Causes very rarely coincident, and to us as yet unknown, be capable of producing a small Degree of Light, either from the greater Density of the Matter, or perhaps from the greater Velocity of its Motion; after the same Manner, as we see the *Effluvia* of *Electric* Bodies emit Light in the Dark.

Monfieur de Mairan has given us a Physical and Historical Treatise of the *Aurora Borealis*, and endeavours to prove that it is owing to the *Zodiacal Light*, or the Atmosphere of the Sun, spread on each Side of it along the *Zodiac* in the form of a *Pyramid*. This, he says, is sometimes extended to such a Length as to reach beyond the Orbit of our Earth, and then mixing itself with our Atmosphere, and being of an *Heterogeneous* Nature, produces

* A round Magnet, so called from the Resemblance it bears to the Earth.

Diff. VII. Of *the* Aurora Borealis. 161

the several Appearances, which are observed in the *Aurora Borealis* *.

I have just mentioned these two Solutions, because they come from two very ingenious Philosophers; though I doubt not but the Reader will agree with me, that they are much too fine spun to hold, and that they are no other than the ingenious *Reveries* of Persons determined to frame an Hypothesis at any Rate. I question not, but we may find Materials for the *Aurora Borealis*, without going so far for them, as these Gentlemen have done; and in particular that we have no Occasion to have Recourse either to the *magnetic Effluvia* of the Earth, or the *Zodiacal Light*, the Nature of both which we are wholly unacquainted with. The Materials employed in the first Solution (I mean such *Effluvia* as are continually exhaled from the Surface and Bowels of the Earth) if rightly considered, will afford a very easy and natural one, as I shall endeavour to shew in the following Manner.

First, we are assured by Experiment, that there are some Steams (as inflammable sulphureous ones) which are capable of so great a Degree of Expansion, that they will render themselves lighter than the Air they float in, though it

* See his Account at large, referred to at the End of Note, Page 155.

be as rare, as it can be made by Art; for they will rise to the Top of the Receiver, when exhausted *, that is when as much Air, as is possible, is drawn out †: Such Steams therefore or Exhalations, rising out of the Earth from Mines, Vulcano's, &c. must necessarily be buoyed up towards the Top of the Atmosphere, at least, till they come to a Region, where the Air is as rare and expanded, as it can be made by the *Air Pump*, here below. This Height according to Dr. *Halley's* Computation §, (which he found upon the Degree of the Air's Elasticity) is about forty or fifty Miles: but probably it is much greater; for the Air, which is higher than Vapours and other *Heterogeneous* Matter that is not elastic, rise to, being much purer than any we can make Experiments upon, may be indued with a much greater Degree of Elasticity, and so the Atmosphere may be considerably higher, than what he upon that Principle, computes it to be.

Secondly, These *Effluvia* being raised to the Top of the Atmosphere, or near it, and floating there, will necessarily be carried towards the poplar Parts thereof, for the following Reasons. 1. Because the superior Current

* See Philosoph. Transact. N^o. 347 and 360.

† It is impossible to extract all the Air out of a Vessel, because it is by the Spring of the Air remaining in the Vessel, that the Valves of the Pump are opened at each Stroke.

§ Philosoph. Transact. N^o. 181,

Diff. VII. *Of the Aurora Borealis.* 163

of the Air, to a great Distance from the *Equator*, is that Way *. 2. We know from Experiment, that whatever swims upon a Fluid which revolves about an *Axis*, is thereby carried towards that *Axis*. This is exactly the Case of these *Effluvia*, for they swim near the Top of the Atmosphere which continually revolves about the *Axis* of the Earth; they must therefore necessarily be carried towards the polar Parts thereof.

Thirdly, These *Effluvia* being collected together at, or near the Poles, and of an inflammable Nature, may easily be supposed to ferment, when they meet with other *heterogeneous* ones proper to produce such an Effect, and emit Streams of Fire; which Streams will naturally rise up into such Parts of the Atmosphere as are still lighter than that wherein the *Effluvia* rest, that is, directly upwards from the Center of the Earth. But, according to the Rules of *Perspective*, those Streams, though they really diverge, as *Radii* from a Center, will appear to a Spectator on the Surface of the Earth to converge towards a Point: Which Point will seem to be directly over his Head, if the Streams ascend in right Lines from the Center of the Earth: but if they deviate all one Way from that Direction, the Point will be on

* As explained in Dissertation V.

164 *Of the Aurora Borealis. Part II.*

that Side the *Zenith* towards which they incline*.

To illustrate this; suppose several Strings hung down from the Ceiling of a Room, and a Candle placed upon a Table below them, the Shadows of them all will converge towards the Point, that is over the Candle. And, if they are made to incline, suppose all one Way, the Point of Convergency will remove from over the Candle, towards that Side of the Room to which the upper Ends of the Strings incline: Now if a Person had viewed them from the Place where the Candle was, and referred their Places to the Ceiling, they would have seemed to him to have converged towards the Point, where their Shadows did.

And if the Streams spread themselves as they arise, but not too much, they will nevertheless appear tapering towards the upper Ends, like *Cones* or *Pyramids*; just as the Sides of a long Walk seem to a Person that views them

* This may be made to appear in the following Manner; Let ADB (Fig. 38.) represent the *Concave* of the Heavens, AB the *Horizon*, C the Place of the Spectator, TV a luminous Substance sending forth the parallel Streams EG, LM, NO, &c. These Streams will all seem to converge towards the Point D, if that Point be taken such, that the Line DC drawn from thence to the Spectator's Eye, be parallel to the Streams. For Instance, the Stream EG will seem to rise from *e* to *g*, LM from *l* to *m*, and FH from *f* to *h*, and so of the rest. And NO will appear wholly in D the Place to which the rest seem to converge. And if the Streams are as large, or somewhat larger at the upper Ends, than at the lower, they will still appear less there, those Ends being farthest from the Spectator's Eye.

from

Diff. VII. Of the Aurora Borealis. 165

from one End of it, or from a distant Place, to approach each other at that which is farthest from him.

This being premised, we may now account for the several *Phænomena* of the *Aurora Borealis* before laid down. As,

1. The Blackness of the Sky, which generally appears in the northern Parts of the *Horizon*, like a dark Cloud, is occasioned by the Brightness of the luminous Matter of the *Aurora* just above it. That the Sky is clear here, is evident (as was observed before) because the smallest Stars are seen through it.

2. The lucid Arch immediately above, is the luminous Matter of the *Aurora* itself, which sometimes exhibits the Appearance of a Curve, sometimes of a strait Line, according to its Form and Situation in the Atmosphere, though generally that of a Curve: For, by the Rules of *Perspective*, when a strait Line is distended horizontally, and above the Spectator's Eye, it ought to appear bent into a Curve, whose Center is below the *Horizon* *. Sometimes it appears on one Side the North Point, more

* Thus, when a Person stands fronting a Row of Houses, and looks over the Tops of them, if they are all of an equal Height, that House which is nearest him, will seem to cut the Heavens in a Point that will be higher than where it is cut by any of the rest; and the Points where the Heavens will seem to be cut by the Tops of those, which are on the Right and Left Hand of the Spectator, will descend lower and lower, as the Houses are farther off; so that the Points, taken all together, will represent a Curve.

than on the other; sometimes regular, sometimes irregular, as the various Circumstances of the Air's Motion at the Top of the Atmosphere, and of the Situation of the flaming Matter may be.

3. The Streams of Light, issuing out of the lucid Arch, are Streams of Fire emitted upwards from the Matter of the *Aurora*, and seem, for the Reasons already laid down, to converge towards the *Zenith* of the Spectator. Why they incline a little sometimes from the Perpendicular, will be explained in the fifth Remark, where we account for the Situation of the Canopy. When no luminous Arch appears, it is probable, that it is intercepted by the *Horizon*, or by the Vapours which float in great Quantities therein.

4. The trembling observed in the upper Part of the Heavens, is owing to the Quickness wherewith the Flashes succeed one another, and also to the irregular Motions and Agitations of the superior Parts of the Atmosphere.

5. So long as the luminous Matter of the *Aurora* is all of it towards the North of us, the Streams cannot seem to meet in a Point at the Top, as will appear to any one that considers the Figure referred to in the Note (p. 164), but after it has advanced forwards, or become kindled over our Heads, then they appear to meet, and form the Canopy already described; and when it has passed further still, they seem
to

Diff.VII. *Of the Aurora Borealis.* 167

to arise from all Parts; though they are much fainter on the southern than on the northern Side, so long as the main Body of the *Aurora* remains on the northern Side of the Canopy, which it rarely, if ever, passes. The Reason why the Center of the Canopy is generally a few Degrees to the South of the Spectator's *Zenith* *, is because the luminous Streams, which issue forth from the extreme Parts of the Substance of the *Aurora*, will naturally diverge a little from the middle ones; and, as those which appear to us, proceed chiefly from the southern Side (that being nearest to us) the Point of Convergency will necessarily be placed to the South of our *Zenith*, according to what was said above about the Inclination of the Strings hanging from the Ceiling of a Room. If the Center of the Canopy is sometimes to the Eastward, and sometimes to the Westward of the Meridian, that depends upon the Motion of that Part of the Air, which is above the Substance of the *Aurora*, and through which the Streams pass, as they rise. This also it is that makes the Streams seem to arise sometimes a little obliquely.

* According to this Theory, the Center of the Canopy will always be near the Spectator's *Zenith*, where-ever he is; which I believe is the Case, for I have met with no Account where it is otherwise; and so every Spectator sees a different Canopy, just as when several Persons are viewing a Rainbow, no two Persons see the same Rainbow at the same Time.

168 *Of the Aurora Borealis. Part II.*

6. The great Height of the *Aurora* is owing to the exceeding Lightness of the *Effluvia*, which compose the Substance of it (as explained above) and the darting of the Streams upwards, into Regions perhaps quite above the Atmosphere, occasions it to exhibit at very distant Places the same Appearances at the same Time.

7. That the *Aurora* appears near the *Pole*, and never at or near the *Equator*, is because of the Tendency the Matter of it has towards the Poles, as explained above. And that it appears in Places more distant from the *Pole*, than it formerly did, is because the *Effluvia*, which are now raised from the Earth, are prevented from approaching so near the polar Parts of the Atmosphere, as they used to do; those Parts being already stocked with others, which were formerly raised, and are now grown *effete* by frequent Fermentations and Explosions.

8. The *horizontal* Trains of Light are the Substance of the *Aurora* just taking Fire, which runs from one Part to another, as in a Train of Gunpowder kindled in any one Part; and sends up Streams perpendicularly from Places, where it meets with a greater Quantity of Matter than ordinary.

9. When the Matter of the *Aurora* is so far spent, as to emit no more Streams, it appears only as a bright steady Light in the North,
which

Diff.VII. *Of the Aurora Borealis.* 169

which gradually dies away, for Want of fresh Fewel to support it.

10. As the Vapours, of which Clouds are formed, never rise so high, as where the Matter of the *Aurora Borealis* floats; it is not at all inconsistent with the foregoing Theory, if it is sometimes intercepted from our Sight, by the Interposition of Clouds below.

See farther on this Subject, *Aristotel.* Meteor. Lib. I. Cap. 4, 5. *Plinii* Histor. Natural. Cap. 26, 27. *Senec.* Quæst. Natural. Lib. I. *Lycost.* Prodigiorum ac Ostentorum Chronicon, passim. *Julius Obsequens* de Prodigiiis, Cap. 13, 43, 88. *Gassendi* Animadvers. in *Diog. Laert.* Lib. X. p. 1157. *Cornelius Gemma* de divinis Naturæ Characterismis. *Nicephori* Histor. Ecclesiast. Lib. XII. Cap. 37. *Isid.* *Hispal.* Histor. Goth. Tom. I. p. 65. Bibliothec. Orientalis Clementino-Vaticana, Tom. I. p. 407. *Gregor. Tur.* passim. Mem. de Lit. de l'Acad. des Inscriptions & belles Lettres, Tom. IV. p. 431. Miscellan. *Berolin.* Tom. I. p. 137. Theatr. Comet. *Stanisf. Lubienietz,* p. 264, 348. Mem. pour servir à l'Histor. de France, Tom. I. p. 168. Mem. de l'Acad. Royal. de Sciences, for almost each Year since 1716. *Philosoph. Transf.* N^o. 305, 310, 320, 347, 348, 349, 351, 352, 363, 365, 368, 376, 385, 395, 398, 399, 402, 410, 418, 431; and the Authors referred to by Mr. *Johnson*, in his Quæst. *Philosoph.* Cap. IV. §. 3, D I S-

DISSERTATION VIII.

Of Fermentation.

HAVING had Occasion to mention some of the Effects of Fermentation, it may not be amiss, before I put an End to these Dissertations, to add a short Account of the Nature of it, and to shew how those Effects are produced by it.

Fermentation is a mutual Commotion of the constituent Particles of Bodies, one among another; and arises from an Inequality in their Attractions of Cohesion. Authors distinguish it into two Kinds; the one is that which happens when a Solid is dissolved by a Fluid; the other is, when two Fluids, being mixed together, ferment with each other.

Those Authors, who have treated of the first of these, tell us, That to cause a Fermentation between a Solid and a Fluid, several Circumstances are necessary. Particularly Dr. *Friend**, and *Keil*† are of Opinion,

1. That the Particles of the Solid must attract those of the Fluid with a greater Force, than the Particles of the Fluid attract one another.

* See his Chemical Lectures.

† See his Letter to Dr. Cockburn, *De Legibus Attractionis*.

2. That

2. That the Pores of the Solid must not be too small to admit the Particles of the Fluid into them.

3. That the Body be of so loose a Contexture, that the Force of Impact, with which the Particles of the Fluid rush into its Pores, may be sufficient to disunite its Parts.

4. That the Elasticity of the Particles tends very much to promote, and augment the Fermentation.

Pr. *Boerhaave* makes also four Conditions requisite *.

1. That there be a due Proportion between the Size of the Particles of the Fluid, and the Pores of the Body to be dissolved.

2. That the Figure of the Particles of the Fluid have a determinate Relation to that of the Pores of the Solid.

3. That the Particles of the Fluid be sufficiently solid, that their Moment, or Force of Action may not be too weak.

4. The last Qualification, he mentions, is a fit Disposition of the Particles of the Fluid, when received into the Pores of the Solid, to make some stay there, and not immediately to pass through; but to act every Way upon the Solid, as they move towards the external Surface thereof.

* Pr. *Boerhaave's* Chemistry, by *Shaw*, p. 344.

But we have no Occasion to have Recourse to so many Suppositions; if the Particles of the Solid attract those of the Fluid with a greater Degree of Force than either those of the Fluid, or those of the Solid attract one another *, it is sufficient; and there will follow a Dissolution of the Body, as may clearly be demonstrated from the Laws of *Mechanics*, whatever the other Circumstances relating to the Figure or Magnitude of Pores, &c. may be †.

When

* This may be thought an impossible Supposition, for the Force of Attraction of Cohesion being as the Surfaces of the attracting Particles, whatever Size or Form the Particles of the Solid and Fluid are of, there cannot be a greater Quantity of Surface between every two Particles, one of which is a Particle of the Solid, and the other a Particle of the Fluid, than there is between every two Particles, which are either both of the Solid, or both of the Fluid; and therefore the Particles of the Solid cannot attract those of the Fluid with greater Force than either those of the Solid or those of the Fluid attract one another. But it is to be considered, that we are not so well acquainted with the Nature of the Attraction of Cohesion, as to determine *exactly* in what Manner, and by what Laws it acts. The Experiments made Use of for this Purpose, only shew that so long as we try them with the same Kind of Bodies, the Attraction is larger where the Contact is so. See Part I. Chap. III. But we have no Method of determining, whether the Difference of Attraction, which *various* Bodies exert upon one another, arises solely from a Difference in their Surfaces, or not.

† *Dem.* Thus, let *f, f, f, &c.* (*Fig. 39.*) represent a Series of the Particles of a Fluid, and *s, s, s, &c.* a Series of those of a solid Body, contiguous to one another: and let the pricked Lines *ff, ff, &c.* represent the Forces of Attraction between the Fluid Particles one among another, and *ss, ss, &c.* those of the solid ones among themselves; and let the black Lines *fs, fs, &c.* express those which are between the fluid and solid Particles. Now, the latter Forces being by the Supposition stronger than the former, the fluid Particles will recede from

Diff.VIII. Of Fermentation. 173

When a Solid is put into a Fluid, if their Particles have the above-mentioned Relation to each other, those of the Solid, being attracted with greater Force towards the Fluid, than they are the contrary Way, they will fall off from the Fluid, and enter in between the Particles of the Fluid; and for the like Reason, those of the Fluid will open to themselves a Way in between those of the Solid, and will separate them from each other. Neither will their Respective Motions cease, unless their

from each other, and suffer those of the Solid to enter in between them; and for the same Reason the solid Particles will give Way to those of the Fluid. By which means, the Distances represented by the pricked Lines becoming greater, the Attraction^s, which they express, will be diminished; so that the fluid Particles will enter quite in between the solid ones, and the solid ones between the fluid one; and both of them together will constitute such a Series, as is represented in *Figure 40*, in the middle Row *s, f, s, f, &c.* where the solid and fluid Particles lie mixed interchangeably one with another in a right Line. Now let it be supposed, that this Series is contiguous to one which consists wholly of Fluid above it, as is expressed in the Figure, and to another below, consisting of solid Particles only. Every solid Particle in this Series will be attracted upwards with greater Force, than it is downwards; and every fluid one with greater Force downwards than it is upwards, as appears by bare Inspection of the Figure, where the black Lines, as in the former, express the stronger Attractions, and the pricked ones the weaker. And, if we suppose the Number of Particles in the Solid and in the Fluid to be nearly equal, those of the Fluid will not stop, till they have quite passed through the Solid; for they will always find a Series wholly consisting of solid Particles before them, whilst that which they leave behind, will be a Mixture of both. In like manner, the solid ones will pass quite through the fluid ones; for they will always meet with more fluid ones before them, than they leave behind within the Sphere of their own Attraction.

Quantities be very unequal, till they are diffused uniformly one among another, as we may very easily conceive; for till then, there will always be some Particles attracted with a greater Degree of Force one Way than they are another.

And if more of the Solid be added to this Fluid, the Particles of the Fluid will also enter into that Solid, till each is surrounded on all Sides with solid Particles, as far as its attractive Force reaches. After which the Fluid is said to be *saturated*, and will dissolve no more.

Again, if more of the Fluid be poured upon that Solid, the solid Particles will diffuse themselves farther into the Fluid, till each of them is encompassed with Particles of the Fluid, as far as its attractive Force extends; and then they will spread themselves no farther.

But in either Case, if another Solid, or Fluid, the attractive Force of whose Particles differ from those of the former, be added, a fresh Fermentation will begin, provided the attractive Forces between the Particles of the former Mixture, and of those which are now added, have such Relation to each other, as is necessary to produce it.

Upon this Principle it should seem, that a Fluid should always be capable of dissolving more than an equal Quantity of a Solid; and that a Solid should be capable of entring in
and

and diffusing itself through more than an equal Quantity of Fluid. The Reason why it is frequently not so, is, because it commonly happens, that the Fluid and the Solid are not of equal specific Gravities. When the Solid is heaviest, so many of its Particles will not ascend and enter into the Fluid, as would otherwise have done; and on the contrary, when the Fluid is heaviest, the Weight of its Particles will be an Impediment to their rising into, and dissolving the Solid.

We have no Occasion to distinguish Fermentation into two Kinds, with regard to its Causes; for, according to the foregoing Theory, whenever two Fluids, or a Solid and a Fluid, are put together, if the Particles of the one attract those of the other, with greater Force than either those of the one, or those of the other attract themselves, a Fermentation will *equally* ensue, the Cause being the same in both Cases.

When two Fluids, or a Solid and a Fluid, ferment with each other, if the Agitation and *intestine* Motion of their Particles be very great, or continues a long Time, and if the Substance of them be of the inflammable Kind, they will, by continually rubbing one against another, be sufficiently heated to take Fire, and burst out into Flame; as was said of the several Compositions mentioned in the foregoing Dissertation.

See

See the Authors, who have explained and defended the old Solution, referred to by Mr. *Johnson* in his *Quæstiones Philosoph.* Cap. III. Quæst. 5, 6, 7.

The End of the Second Part.



II.

nd

Ir.

II.

P. 105. *P.*

110.




Fig. 37. P. 135.




Fig. 39. P. 172.




Fig. 40. P. 173.




Fig. 38. P. 164.

